



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 15/12, C07K 14/71, 14/51, 16/28, A61K 38/18</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 95/07982</b> <b>(43) International Publication Date:</b> 23 March 1995 (23.03.95)
<b>(21) International Application Number:</b> PCT/US94/10080 <b>(22) International Filing Date:</b> 7 September 1994 (07.09.94)  <b>(30) Priority Data:</b> 08/123,934 17 September 1993 (17.09.93) US  <b>(71) Applicant:</b> GENETICS INSTITUTE, INC. [US/US]; 87 CambridgePark Drive, Cambridge, MA 02140 (US).  <b>(72) Inventors:</b> WOZNEY, John, M.; 59 Old Bolton Road, Hudson, MA 01749 (US). CELESTE, Anthony, J.; 86 Packard Street, Hudson, MA 01749 (US). THIES, R., Scott; 10 McKenney Circle, Andover, MA 01810 (US). YAMAJI, Noboru; 21 Miyukigaoka, Tsukuba-City, Ibaraki 305 (JP).  <b>(74) Agent:</b> LAZAR, Steven, R.; Genetics Institute, Inc., 87 CambridgePark Drive, Cambridge, MA 02140 (US).		<b>(81) Designated States:</b> AU, CA, FI, JP, KP, KR, NO, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>  <b>BEST AVAILABLE COPY</b>
<b>(54) Title:</b> ACTIVIN RECEPTORS-LIKE KINASE (ALK), BELONGING TO THE TGF RECEPTOR FAMILY AND/OR TO THE BMP RECEPTOR FAMILY  <b>(57) Abstract</b>  Novel serine/threonine receptor proteins and BMP receptor proteins are disclosed, as well as DNA molecules encoding said proteins and methods of using the receptor proteins. Further disclosed are truncated BMP receptor proteins and molecules which act as ligands to said BMP receptor proteins.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgystan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

# ACTIVIN RECEPTORS-LIKE KINASE (ALK), BELONGING TO THE TGF RECEPTOR FAMILY AND/OR TO THE BMP RECEPTOR FAMILY

## FIELD OF THE INVENTION

The present invention relates to novel serine/threonine kinase receptor proteins, including a novel family of receptor proteins to bone morphogenetic proteins (BMPs). More particularly, the present invention relates to receptor proteins which are able to bind to BMPs, including BMP-2 and BMP-4. The present invention further relates to methods of isolating novel BMP receptor proteins using newly identified DNA fragments as probes for isolating such proteins.

## BACKGROUND OF THE INVENTION

Bone morphogenetic proteins (BMPs) are a family of proteins which have been identified as having the ability to induce the formation of bone and cartilage in tissue extracts. BMPs are a subfamily within the TGF- $\beta$  superfamily. BMPs have multiple therapeutic uses, including a wide variety of settings where bone has been lost through physiological or traumatic processes.

The TGF- $\beta$  superfamily of proteins have been shown to bind to serine/threonine kinase receptors. Massague, Cell, 69:1067-1070 (1992); Attisano et al., Cell 68:97-108 (1992); Lin et al., Cell, 68:775-785 (1992); Wang et al., Cell 67:797-805 (1991). Similarly, activin receptors have been isolated and characterized as a predicted transmembrane serine kinase. Mathews et al., Cell 65:973-982 (1991); Nakamura et al., J. Biol. Chem. 267:18924-18928 (1992). Ebner et al., Science, 260:1344-1348 (1993) describe the existence of Type I and Type II TGF- $\beta$  receptors, and the effects of the Type I receptor on binding of TGF- $\beta$  to the Type II receptor.

Type I receptor proteins have been reported not to bind to their ligand molecules independently, but, acting in concert with Type II receptor proteins, are observed to contribute to increased binding to the ligand. See Matsuzaki et al., J. Biol. Chem., 268:12719-12723 (1993); Ebner et al., Science, 260:1344-1348 (1993).

Paralkar et al., PNAS USA 88:3397-3401 (1991) describes the presence of high affinity binding sites for BMP-4 on MC3T3E1 and NIH3T3 cells. No competition by TGF- $\beta$  was found for the BMP-4 binding proteins, nor was competition by BMP-4 for TGF- $\beta$  receptors observed in Attisano et al., Cell 68:97-108 (1992).

## SUMMARY OF THE INVENTION

In one embodiment, the present invention comprises a purified and isolated DNA molecule which encodes a BMP receptor protein, said DNA molecule preferably comprising the clones CFK1-43a and CFK1-23a, or a DNA sequence selected from the group consisting of SEQ ID NO:1 and SEQ ID NO:3.

The present invention further comprises purified and isolated DNA molecules which encode BMP receptor proteins, said BMP receptor proteins preferably comprising an amino acid sequence selected from the group consisting of SEQ ID NO:2 and SEQ ID NO:4. In another embodiment, the present invention comprises a BMP receptor protein CFK1-43a and CFK1-23a, comprising an amino acid sequence selected from the group consisting of SEQ ID NO:2 and SEQ ID NO:4.

The present invention further comprises DNA molecules comprising a DNA sequence selected from the group consisting of SEQ ID NO:5, SEQ ID NO:7 and SEQ ID NO:9, and DNA molecules which encode serine/threonine kinase receptor proteins comprising an amino acid sequence selected from the group consisting of SEQ ID NO:6, SEQ ID NO:8 and SEQ ID NO:10. These DNA molecules and proteins are related to the BMP family of receptors. Among other uses, these DNA molecules are presently useful as probes for isolating and purifying additional novel BMP receptors.

The present invention also comprises novel DNA sequences which encode receptor proteins, which novel DNA sequences are identified by a method using DNA sequence encoding all or a fragment of the receptor proteins of the present invention. In preferred embodiments, the novel DNA sequences are identified using DNA sequence from the serine/threonine kinase domain of a receptor, which is highly conserved among the family of BMP receptors. Alternatively, DNA sequence encoding the ligand binding domain could be used to identify additional novel BMP receptor encoding sequences.

The present invention further comprises DNA molecules encoding soluble, truncated receptor proteins, and the soluble proteins themselves. The truncated receptor proteins preferably comprise the ligand binding domain, but not the serine/threonine kinase and transmembrane domains, of the receptor protein. The truncated receptor proteins are soluble, and will be secreted into supernatant by mammalian cells. Thus, when expressed



in mammalian cells using a DNA molecule encoding a truncated receptor protein, the truncated receptor protein will be secreted rather than expressed on the surface of the host cell. The truncated receptor protein thereby expressed still binds specifically to BMPs, and can be used to block receptors from mediating the cellular processes in which they normally participate in as signalling mechanisms by competition for the same ligand. The truncated receptor protein could compete with receptor proteins normally expressed on the surface of responsive cells for functional ligand and inhibit the formation of a functional receptor-ligand complex, thereby blocking the normal signalling mechanism of the complex and the cellular processes normally affected by functional receptor-ligand interactions.

10 In one aspect, the invention provides a method for producing cells expressing more than one receptor protein comprising culturing a selected host cell containing a polynucleotide sequence encoding a first selected receptor protein, truncated receptor protein, or active fragment thereof and a polynucleotide sequence encoding a second selected receptor protein, truncated receptor protein, or active fragment thereof. The  
15 resulting cells, which will express multiple co-expressed, biologically active receptors, may be isolated and used in a therapeutic composition.

Another aspect of the current invention comprises ligands for the BMP receptors and truncated BMP receptor protein, said ligands being characterized by the ability to bind to the receptors. Such ligands may stimulate growth of bone and/or cartilage, or may be  
20 involved in influencing other developmental processes. Said ligands may be monoclonal antibodies, small peptide BMP analogues, or small organic molecule BMP analogues as further characterized herein. In a preferred embodiment, said ligands comprise antibodies against the truncated, soluble receptor protein and the receptor proteins of the invention. These antibodies can be employed in a variety of diagnostic and therapeutic applications.  
25 Such antibodies can be used to identify cell types which naturally express receptors of the invention and may therefore have the capacity to elicit a biological response upon exposure to the appropriate ligand. These antibodies can be further useful in the identification of additional receptor proteins capable of binding to other individual BMPs and/or BMP heterodimers. Additionally such antibodies are useful in blocking the formation of  
30 functional receptor-ligand complexes and thus inhibit the cellular responses that would

normally be mediated by these complexes. Alternatively, such antibodies may mimic the effect of BMP by interacting with the receptor in a way that would stimulate the cellular responses that would normally be mediated by a functional receptor-ligand complex.

5 In yet another embodiment, the invention comprises pharmaceutical compositions comprising a compound first identified for such use as a ligand for the truncated BMP receptor and therapeutic methods for the treatment of bone and/or cartilage disorders comprising administering a ligand for the truncated BMP receptor.

Other aspects and advantages of the present invention will be apparent upon consideration of the following detailed description and preferred embodiments thereof.

10 Brief Description of the Sequences

SEQ ID NO:1 comprises DNA and amino acid sequence of the BMP receptor protein CFK1-23a, isolated from rat cell line CFK1. This DNA contained in plasmid CFK1-23a, which has been deposited and accorded ATCC #69378, further described below.

15 SEQ ID NO:2 comprises the amino acid sequence encoded by the CFK1-23a DNA sequence.

SEQ ID NO:3 comprises DNA and amino acid sequence of the BMP receptor protein CFK1-43a, isolated from rat cell line CFK1. This DNA contained in plasmid CFK1-43a has been deposited and accorded ATCC #69381, further described below.

20 SEQ ID NO:4 comprises the amino acid sequence encoded by the CFK1-43a DNA sequence.

SEQ ID NO:5 comprises DNA and amino acid sequence of the serine/threonine kinase receptor protein CFK1-10a, isolated from rat cell line CFK1. This DNA contained within plasmid CFK1-10a has been deposited and accorded ATCC #69380, further described below.

25 SEQ ID NO:6 comprises the amino acid sequence encoded by the CFK1-10a DNA sequence.

SEQ ID NO:7 comprises DNA and amino acid sequence of the serine/kinase receptor protein W101, isolated from murine cell line W-20-17. This DNA contained in plasmid pMT101 has been deposited and accorded ATCC #69379, further described

30

below.

SEQ ID NO:8 comprises the amino acid sequence encoded by the W101 DNA sequence.

5 SEQ ID NO:9 comprises DNA and amino acid sequence of the serine/kinase receptor protein W120, isolated from murine cell line W-20-17. This DNA contained in plasmid pMT120E has been deposited and accorded ATCC #69377, further described below.

SEQ ID NO:10 comprises the amino acid sequence encoded by the W120 DNA sequence.

10 SEQ ID NO:11 comprises DNA and amino acid sequence of the serine/kinase receptor protein KDA-B5. This DNA was used as a probe to identify novel serine/kinase receptors of the present invention.

SEQ ID NO:12 comprises the amino acid sequence encoded by the KDA-B5 DNA sequence.

15 SEQ ID NO:13: comprises the DNA sequence of oligonucleotide primer A.

SEQ ID NO:14: comprises the DNA sequence of oligonucleotide primer B.

SEQ ID NO:15: comprises the DNA sequence of oligonucleotide primer C..

SEQ ID NO:16 comprises the DNA sequence of oligonucleotide primer D.

SEQ ID NO:17 comprises the DNA sequence of oligonucleotide primer E.

20 SEQ ID NO:18: comprises the amino acid sequence of a portion of KDA-B5 used to design oligonucleotide primer A.

SEQ ID NO:19 comprises the amino acid sequence of a portion of KDA-B5 used to design oligonucleotide primer B through E.

#### Detailed Description of the Invention

25 Bone morphogenetic proteins are characterized by their ability to promote, stimulate or otherwise induce the formation of cartilage and/or bone. The ability of these proteins to demonstrate cartilage and/or bone formation activity in the rat bone formation assay described below. These proteins can be used in compositions which may be used to induce bone and/or cartilage formation. These BMP compositions may also be used for  
30 wound healing and tissue repair. Further uses of such compositions include the treatment

of bone and/or cartilage defects, periodontal disease and other tooth repair processes, treatment of osteoporosis and increase of neuronal survival.

5 The BMP receptors and truncated receptors of the present invention are useful, among other uses, for the identification of BMPs, the identification of further BMP receptors, and the identification of ligands or molecules, including antibodies, which are able to mimic the binding characteristics of BMPs. These ligands may act as agonist or antagonists, depending upon the individual ligand. The activity of the ligands may be characterized in an assay for BMP activity, such as the W-20-17 alkaline phosphatase induction assay and rat ectopic bone formation assay, described at Examples XII and XIII below. The BMP receptors are also useful in inhibiting the effects of BMPs, where such inhibition is desired.

BMP receptor proteins of the present invention may be characterized by an amino acid sequence comprising amino acid # 1-532 of SEQ ID NO:2; or amino acid # 1-502 of SEQ ID NO:4.

15 The purified human BMP receptor proteins of the present invention may be produced by culturing a host cell transformed with a DNA sequence comprising the DNA coding sequence of SEQ ID NO:1 from nucleotide # 61 to nucleotide 1656 (or to 1659 with the stop codon); or SEQ ID NO:3 from nucleotide #247 to nucleotide 1752 (or to 1755 stop codon); and recovering and purifying from the transformed cell membrane a protein which contains the derived amino acid sequence, or a substantially homologous sequence as represented by amino acid # 24 to # 532 of SEQ ID NO:2; or amino acid # 8 to # 502 of SEQ ID NO:4. Since the BMP receptor proteins expressed in this manner are expected to remain associated with the cell membrane of the transformed cell, recombinant receptor proteins of the invention can be dissociated from the transformed cell membrane and are then purified by isolating them from other proteinaceous materials with which they are co-produced and from other contaminants present.

25 Truncated BMP receptor proteins of the present invention may be characterized by an amino acid sequence comprising amino acid # 1-149 of SEQ ID NO:2; or amino acid # 1-124 of SEQ ID NO:4.

30 The purified human truncated BMP receptor proteins of the present invention may

be produced by culturing a host cell transformed with a DNA sequence comprising the DNA coding sequence of SEQ ID NO:1 from nucleotide # 61 to nucleotide 507; or SEQ ID NO:3 from nucleotide # 247 to nucleotide 618; and recovering and purifying from the culture medium a protein which contains the derived amino acid sequence, or a substantially homologous sequence, as represented by amino acid # 24 to # 149 of SEQ ID NO:2; or amino acid # 8 to # 124 of SEQ ID NO:4. In the above amino acid sequences, the secretory leader sequence (e.g., amino acids 1 to 23 of SEQ ID NO:2) will not be present since these are typically cleaved away from secreted proteins. The leader sequence predicted for SEQ ID NO:4 by standard computer programs is amino acids 1 to 7; however, it is contemplated that the actual leader sequence may be longer since seven amino acids is unusually short for a leader sequence. Thus, the protein purified from culturing host cells transformed with a DNA molecule comprising the DNA sequence of SEQ ID NO:3 from nucleotide # 247 to nucleotide 618 may be shorter than amino acid # 8 to # 124 of SEQ ID NO:4.

The truncated BMP receptor proteins recovered from the culture medium are purified by isolating them from other proteinaceous materials with which they are co-produced and from other contaminants present.

Other serine/threonine kinase receptor proteins of the present invention may be characterized by an amino acid sequence comprising amino acid # 1-509 of SEQ ID NO:6.

The purified serine/threonine kinase receptor proteins of the present invention may be produced by culturing a host cell transformed with a DNA sequence comprising the DNA coding sequence of SEQ ID NO:5 from nucleotide # 474 to nucleotide 2000 (or to 2003 stop codon); and recovering and purifying from the transformed cell membrane a protein which contains the derived amino acid sequence, or a substantially homologous sequence as represented by amino acid # 18 to # 509 of SEQ ID NO:6. Since the serine/threonine kinase receptor proteins expressed in this manner are expected to remain associated with the cell membrane of the transformed cell, recombinant receptor proteins of the invention can be dissociated from the transformed cell membrane and are then purified by isolating them from other proteinaceous materials with which they are co-produced and from other contaminants present.



Truncated serine/threonine kinase receptor proteins of the present invention may be characterized by an amino acid sequence comprising amino acid # 1-121 of SEQ ID NO:6.

5 The purified human truncated serine/threonine kinase receptor proteins of the present invention may be produced by culturing a host cell transformed with a DNA sequence comprising the DNA coding sequence of SEQ ID NO:5 from nucleotide # 474 to nucleotide 836 and recovering and purifying from the culture medium a protein which contains the derived amino acid sequence, or a substantially homologous sequence as represented by amino acid # 18 to # 121 of SEQ ID NO:6. The truncated serine/threonine  
10 kinase receptor proteins recovered from the culture medium are purified by isolating them from other proteinaceous materials with which they are co-produced and from other contaminants present.

Serine/kinase receptor proteins of the present invention may be characterized by an amino acid sequence comprising amino acid # 1-505 of SEQ ID NO:8; or amino acid  
15 # 1-503 of SEQ ID NO:10.

The purified serine/kinase receptor proteins of the present invention may be produced by culturing a host cell transformed with a DNA sequence comprising the DNA coding sequence of SEQ ID NO:7 from nucleotide # 80 to nucleotide 1594 (or to 1597 stop codon); or SEQ ID NO:9 from nucleotide # 83 to nucleotide 1591 (or to 1594 stop  
20 codon); and recovering and purifying from the transformed cell membrane a protein which contains the derived amino acid sequence, or a substantially homologous sequence as represented by amino acid # 24 to # 505 of SEQ ID NO:8; or amino acid # 30 to # 503 of SEQ ID NO:10. Since the serine/kinase receptor proteins expressed in this manner are expected to remain associated with the cell membrane of the transformed cell, recombinant  
25 receptor proteins of the invention can be dissociated from the transformed cell membrane and are then purified by isolating them from other proteinaceous materials with which they are co-produced and from other contaminants present.

Truncated serine/threonine kinase receptor proteins of the present invention may be characterized by an amino acid sequence comprising amino acid # 1-122 of SEQ ID  
30 NO:8; or amino acid # 1-121 of SEQ ID NO:10.



The purified human truncated serine/threonine kinase receptor proteins of the present invention may be produced by culturing a host cell transformed with a DNA sequence comprising the DNA coding sequence of SEQ ID NO:7 from nucleotide # 80 to nucleotide 445; or SEQ ID NO:9 from nucleotide # 83 to nucleotide 445; and recovering and purifying from the culture medium a protein which contains the derived amino acid sequence, or a substantially homologous sequence, as represented by amino acid # 24 to # 122 of SEQ ID NO:8; or amino acid # 30 to # 121 of SEQ ID NO:10. The truncated serine/threonine kinase receptor proteins recovered from the culture medium are purified by isolating them from other proteinaceous materials with which they are co-produced and from other contaminants present.

The present invention also encompasses DNA molecules comprising the novel DNA sequences, free of association with DNA sequences encoding other proteinaceous materials, and coding for the expression of the above receptor proteins. These DNA sequences include those depicted in SEQ ID NOS: 1, 3, 5, 7 and 9, in a 5' to 3' direction and those sequences which hybridize under stringent hybridization conditions [see, T. Maniatis et al, Molecular Cloning (A Laboratory Manual), Cold Spring Harbor Laboratory (1982), pages 387 to 389] to the DNA sequences of SEQ ID NOS: 1, 3, 5, 7 and 9; and encode a protein having the ability to bind to BMP or which is useful to isolate novel BMP receptors.

Similarly, DNA sequences which code for the above receptor polypeptides coded for by the amino acid sequences of SEQ ID NO: 2, 4, 6, 8 and 10, but which differ in codon sequence due to the degeneracies of the genetic code or allelic variations (naturally-occurring base changes in the species population which may or may not result in an amino acid change) also encode the novel receptor proteins described herein. Variations in the DNA sequences of SEQ ID NOS: 1, 3, 5, 7 and 9 which are caused by point mutations or by induced modifications (including insertion, deletion, and substitution) to enhance the activity, half-life or production of the polypeptides encoded thereby are also encompassed in the invention.

Another aspect of the present invention provides a novel method for producing receptor proteins. The method of the present invention involves culturing a suitable cell

line, which has been transformed with a DNA molecule comprising a DNA sequence coding on expression for a receptor protein, under the control of known regulatory sequences. The transformed host cells are cultured and the receptor proteins recovered and purified from the transformed cell membrane. The purified proteins are substantially free from other proteins with which they are co-produced as well as from other contaminants.

Another aspect of the present invention provides a novel method for producing truncated receptor proteins. The method of the present invention involves culturing a suitable cell line, which has been transformed with a DNA molecule comprising a DNA sequence coding on expression for a truncated receptor protein, under the control of known regulatory sequences. The transformed host cells are cultured and the truncated receptor proteins recovered and purified from the culture medium. The purified proteins are substantially free from other proteins with which they are co-produced as well as from other contaminants.

Suitable cells or cell lines for production of the receptor proteins or truncated receptor proteins may be mammalian cells, such as Chinese hamster ovary cells (CHO) or BHK cells. The selection of suitable mammalian host cells and methods for transformation, culture, amplification, screening and product production and purification are known in the art. See, e.g., Gething and Sambrook, Nature, 293:620-625 (1981), or alternatively, Kaufman et al, Mol. Cell. Biol., 5(7):1750-1759 (1985) or Howley et al, U.S. Patent 4,419,446. Another suitable mammalian cell line, which is described in the accompanying examples, is the monkey COS-1 cell line. The mammalian cell line CV-1 may also be suitable.

Bacterial cells may also be suitable hosts. For example, the various strains of E. coli (e.g., HB101, MC1061) are well-known as host cells in the field of biotechnology. Various strains of B. subtilis, Pseudomonas, other bacilli and the like may also be employed in this method.

Many strains of yeast cells known to those skilled in the art may also be available as host cells for expression of the polypeptides of the present invention. Additionally, where desired, insect cells may be utilized as host cells in the method of the present

invention. See, e.g. Miller et al, Genetic Engineering, 8:277-298 (Plenum Press 1986) and references cited therein.

Another aspect of the present invention provides vectors for use in expression of these novel receptor polypeptides. Preferably, the vectors contain the full novel DNA sequences described above which encode the novel receptor proteins of the invention. Additionally, the vectors contain appropriate expression control sequences permitting expression of the receptor protein sequences.

Alternatively, vectors incorporating modified DNA sequences as described above are also embodiments of the present invention and useful in the production of the receptor proteins. The vectors may be employed in the method of transforming cell lines and contain selected regulatory sequences in operative association with the DNA coding sequences of the invention which are capable of directing the replication and expression thereof in selected host cells. Useful regulatory sequences for such vectors are known to one of skill in the art and may be selected depending upon the selected host cells. Such selection is routine and does not form part of the present invention.

The BMP receptor proteins of the present invention, such as CFK1-23a and CFK1-43a, have been found to bind to members of the BMP family, preferably BMP-2 and BMP-4, but not to TGF- $\beta$ . Thus, the BMP receptor proteins of the present invention are distinguished from TGF- $\beta$  receptors, which bind to TGF- $\beta$ .

The present invention may include co-transfection of cells with DNA molecules comprising DNA sequences encoding multiple receptor proteins in order to achieve binding to a ligand molecule such as a BMP. Thus, for example, a DNA molecule comprising a DNA sequence encoding the receptor protein CFK1-10a may be co-transfected into cells along with a DNA molecule comprising a DNA sequence encoding receptor protein CFK1-23a or CFK1-43a.

The DNA molecules comprising DNA sequences encoding the receptor proteins of the present invention are useful for the production of cells which express receptor proteins. These cells, when transformed with the DNA molecules of the present invention, will express receptor proteins on their surface. In turn, these cells will bind more readily to the ligand and may demonstrate increased responsiveness to the ligand. For example, cells

which express the BMP receptor proteins of the present invention exhibit increased binding to BMP-2 and BMP-4, and will exhibit increased responsiveness to BMPs such as BMP-2 and BMP-4. The increased BMP response is desirable for accelerating the effects of BMPs, which include the osteoinductive promotion of bone growth and cartilage regeneration.

The BMP receptor proteins of the present invention are useful for isolating BMP. Additionally, BMP receptor proteins of the invention are useful in the identification of novel molecules related to BMPs which may be capable of inducing the formation of bone or cartilage or may be involved in influencing other developmental processes. In addition, the BMP receptor proteins are useful for identifying and/or quantifying BMP-2 and/or BMP-4 in a sample, as well as for inhibiting the effects of BMP-2 or BMP-4 on cells. The BMP receptors of the present invention may further be useful in identifying synthetic and naturally-occurring chemical entities which are able to mimic the binding effects of BMP-2 and/or BMP-4. The BMP receptor proteins of the invention may also be useful in identifying synthetic and naturally-occurring chemical entities which are able to antagonize and/or inhibit the binding effects of BMP-2 and/or BMP-4. The BMP receptor proteins may also be useful in identifying compounds which play a role in regulating the expression of BMP receptor proteins. Those compounds could be used in order to stimulate BMP-responsiveness, for example, bone growth, in particular tissues or cells of interest.

The novel serine/threonine kinase receptor proteins of the present invention also include W101 and W120, which have been isolated from murine cell line W-20-17, a cell line which is known to be responsive to BMP. The DNA encoding one of these novel receptor proteins has been used as a probe in order to isolate other clones which are potentially members of the class of BMP receptor proteins, including the CFK1-23a and CFK1-43a clones, which have been confirmed to encode proteins which are members of the BMP receptor family. Thus, the DNA molecules comprising DNA sequence encoding the serine/kinase receptor proteins of the present invention are useful for the isolation of DNA encoding BMP receptor proteins, and the present invention includes such a method of using the DNA molecules comprising DNA sequence encoding serine/threonine kinase



receptor proteins, as well as the novel BMP receptor proteins which are thereby isolated.

In one embodiment of the present invention, novel DNA sequences which encode BMP receptor proteins are identified by a method using DNA sequence encoding all or a fragment of the serine/kinase receptor proteins of the present invention. In preferred  
5       embodiments, the novel DNA sequences are identified using DNA sequence encoding the serine/threonine kinase domain of a receptor. Alternatively, DNA sequence encoding the ligand binding domain could be used to identify additional novel BMP receptor encoding sequences.

Thus, the present invention further comprises methods of identifying new BMP  
10       receptor proteins and DNA molecules encoding those proteins, and the proteins and DNA molecules thus identified. The method comprises preparing a DNA fragment which encodes a selected domain of a BMP receptor protein, preferably the kinase domain of a BMP receptor protein, or alternatively a DNA fragment encoding the ligand binding domain, and using that fragment as a probe to screen either a genomic or cDNA library.  
15       The cDNA library is preferably prepared from a cell line known to express BMP receptors. These include the murine cell line W-20-17 and the rat cell line CFK1. The DNA sequences which are thus identified share homology with the known BMP receptor protein, and thus are expected to encode a protein which will bind to one or more BMPs. Using methods known in the art, one can clone the entire DNA sequence which is thereby  
20       identified and use it to express the newly identified BMP receptor protein. Identification of the new protein as a BMP receptor protein is confirmed using the binding assay described in Example VI.

Another embodiment of the present invention comprises DNA molecules comprising DNA sequences encoding truncated receptor proteins, and the truncated proteins  
25       themselves. The truncated receptor proteins preferably comprise the ligand binding domain, but not the serine/threonine kinase and transmembrane domains, of the receptor protein. The truncated receptor proteins are soluble, and will be secreted into supernatant by mammalian cells. Thus, when expressed in mammalian cells using a DNA molecule encoding a truncated receptor protein, the truncated receptor protein will be secreted rather  
30       than expressed on the surface of the host cell. The truncated receptor protein thereby

expressed still binds specifically to its ligand. Thus, the truncated BMP receptor proteins can be used to block BMP receptors of the invention from mediating the cellular processes in which they normally participate in as signalling mechanisms. The truncated receptor protein could compete with receptor proteins normally expressed on the surface of responsive cells for functional ligand and inhibit the formation of a functional receptor-ligand complex, thereby blocking the normal signalling mechanism of the complex and the cellular processes normally affected by functional receptor-ligand interactions.

Compositions containing the truncated BMP receptor proteins of the present invention may be used for the inhibition of the effects of BMPs such as BMP-2 and/or BMP-4 on cells. The present invention includes therapeutic methods comprising administering such a composition topically, systemically, or locally as an implant or device. When administered, the therapeutic composition for use in this invention is, of course, in a pyrogen-free, physiologically acceptable form. Further, the composition may desirably be encapsulated or injected in a viscous form for delivery to the desired site. Therapeutically useful agents, such as growth factors (e.g., BMPs, TGF- $\beta$ , FGF, IGF), cytokines (e.g., interleukins and CSFs) and antibiotics, may also optionally be included in or administered simultaneously or sequentially with, the receptor composition in the methods of the invention.

Another embodiment of the present invention comprises cells which have been transformed with the DNA molecules comprising DNA sequences encoding the BMP receptor proteins of the present invention. These cells will express BMP receptors on their surface, which will increase the cells' responsiveness to BMP. Thus, cells transformed with the DNA molecules encoding the BMP receptor proteins may be administered therapeutically, to promote response to BMP, for example, bone and/or cartilage regeneration at a desired site.

There is a wide range of methods which can be used to deliver the cells expressing BMP receptor proteins to a site for use in promoting a BMP response such as bone and or cartilage regeneration. In one embodiment of the invention, the cells expressing BMP receptor protein can be delivered by direct application, for example, direct injection of a sample of such cells into the site of bone or cartilage damage. In a particular embodiment,



these cells can be purified. In a preferred embodiment, the cells expressing BMP receptor protein can be delivered in a medium or matrix which partially impedes their mobility so as to localize the cells to a site of bone or cartilage injury. Such a medium or matrix could be semi-solid, such as a paste or gel, including a gel-like polymer. Alternatively, the medium or matrix could be in the form of a solid, preferably, a porous solid which will allow the migration of cells into the solid matrix, and hold them there while allowing proliferation of the cells.

In a method of the present invention, the cells expressing BMP receptors are applied in the desired site as described above, and BMP is applied. The BMP may be applied simultaneously or immediately following application of the cells expressing BMP receptors. BMPs are known and have been described as follows: BMP-2 (sometimes referred to as BMP-2A) and BMP-4 (sometimes referred to as BMP-2B), U.S. Patent No. 5,013,649; BMP-3 U.S. Patent No. 5,116,738; BMP-5, U.S. Patent No. 5,106,748; BMP-6, U.S. Patent No. 5,187,076; BMP-7, U.S. Patent No. 5,141,905; BMP-8, PCT Publication No. WO93/00432; BMP-9, Serial No. 07/720,590, filed on June 25, 1991; BMP-10, Serial No. \_\_\_\_\_, filed on May 12, 1993. Heterodimers are described in United States Patent Application Serial No. 07/787,496, filed on April 7, 1992. The disclosure of the above references are hereby incorporated herein by reference as if fully reproduced herein. The BMP may be applied in manners known in the art, such as described in the above patents, as well as in United States Patent 5,171,579, the disclosure of which is also hereby incorporated by reference.

#### Expression of Receptor Protein

In order to produce receptor protein, the DNA encoding the desired protein is transferred into an appropriate expression vector and introduced into mammalian cells or other preferred eukaryotic or prokaryotic hosts by conventional genetic engineering techniques. The presently preferred expression system for biologically active recombinant receptor protein is stably transformed mammalian cells.

One skilled in the art can construct mammalian expression vectors by employing the sequence of SEQ ID NO: 1, 3, 5, 7 or 9, or other DNA sequences containing the coding sequences of SEQ ID NO: 1, 3, 5, 7 or 9, or other modified sequences and known

vectors, such as pCD [Okayama et al., Mol. Cell Biol., 2:161-170 (1982)] and pJL3, pJL4 [Gough et al., EMBO L., 4:645-653 (1985)]. The receptor protein cDNA sequences can be modified by removing the non-coding nucleotides adjacent to the 5' and 3' ends of the coding region. The deleted non-coding nucleotides may or may not be replaced by other sequences known to be beneficial for expression. The transformation of these vectors into appropriate host cells can result in expression of receptor proteins.

One skilled in the art can manipulate the sequences of SEQ ID NO: 1, 3, 5, 7 or 9 by eliminating or replacing the mammalian regulatory sequences flanking the coding sequence with bacterial sequences to create bacterial vectors for intracellular or extracellular expression by bacterial cells. For example, the coding sequences can be further manipulated (e.g. ligated to other known linkers or modified by deleting non-coding sequences therefrom or altering nucleotides therein by other known techniques). The modified receptor protein coding sequence can then be inserted into a known bacterial vector using procedures such as described in T. Taniguchi et al., Proc. Natl Acad. Sci. USA, 77:5230-5233 (1980). This exemplary bacterial vector can then be transformed into bacterial host cells and receptor protein expressed thereby. For a strategy for producing extracellular expression of receptor proteins in bacterial cells., see, e.g. European patent application EPA 177,343.

Similar manipulations can be performed for the construction of an insect vector [See, e.g. procedures described in published European patent application 155,476] for expression in insect cells. A yeast vector can also be constructed employing yeast regulatory sequences for intracellular or extracellular expression of the receptor proteins of the present invention by yeast cells. [See, e.g., procedures described in published PCT application WO86/00639 and European patent application EPA 123,289].

A method for producing high levels of a receptor protein of the invention in mammalian cells involves the construction of cells containing multiple copies of one or more of the heterologous receptor genes. The heterologous gene is linked to an amplifiable marker, e.g. the dihydrofolate reductase (DHFR) gene for which cells containing increased gene copies can be selected for propagation in increasing concentrations of methotrexate (MTX) according to the procedures of Kaufman and Sharp,

J. Mol. Biol., 159:601-629 (1982). This approach can be employed with a number of different cell types.

For example, a plasmid containing a DNA sequence for a BMP receptor protein of the invention in operative association with other plasmid sequences enabling expression thereof and the DHFR expression plasmid pAdA26SV(A)3 [Kaufman and Sharp, Mol. Cell. Biol., 2:1304 (1982)] can be co-introduced into DHFR-deficient CHO cells, DUKX-BII, by calcium phosphate coprecipitation and transfection, electroporation, protoplast fusion or lipofection. DHFR expressing transformants are selected for growth in alpha media with dialyzed fetal calf serum, and subsequently selected for amplification by growth in increasing concentrations of MTX (e.g. sequential steps in 0.02, 0.2, 1.0 and 5uM MTX) as described in Kaufman et al., Mol Cell Biol., 5:1750 (1983).

Transformants are cloned, and binding to BMP-2 or BMP-4 is measured by the binding assay described above in Example VI. BMP-2 and BMP-4 binding should increase with increasing levels of MTX resistance. Similar procedures can be followed to produce other related BMP receptor proteins.

#### Co-Expression of Multiple Receptor Proteins

According to one embodiment of this invention, the host cell may be co-transfected with one or more vectors containing coding sequences for one or more receptor proteins, truncated receptor proteins or active fragments thereof. Each receptor polynucleotide sequence may be present on the same vector or on individual vectors co-transfected into the cell. Alternatively, the polynucleotides encoding receptors, truncated receptors or their fragments may be incorporated into a chromosome of the host cell. Additionally, a single transcription unit may encode single copy of two genes encoding different receptor proteins.

According to another embodiment of this invention, the selected host cell containing the two polypeptide encoding sequences is a hybrid cell line obtained by fusing two selected, stable host cells, each host cell transfected with, and capable of stably expressing, a polynucleotide sequence encoding a selected first or second receptor protein, truncated receptor protein or active fragment thereof.

In another aspect of the present invention, therefore, there are provided

compositions of cells which express more than one recombinant receptor protein, truncated receptor protein; or active fragments thereof which retain the binding characteristics of the receptor or truncated receptor. Also provided are compositions of truncated truncated receptor proteins secreted by host cells. The cells, proteins, and compositions of receptor proteins, truncated receptor proteins or active fragments thereof may be characterized by their ability to bind selectively to BMPs with greater binding affinity than to other proteins in the TGF- $\beta$  superfamily in a binding assay.

The cells and compositions may comprise one or more BMP receptor proteins, truncated BMP receptor proteins, or active fragments thereof; or of one or more serine/threonine kinase receptor proteins, truncated serine/threonine kinase receptor proteins, or active fragments thereof, such as W-101, W-120 or CFK1-10a, in combination with one or more BMP receptor proteins, truncated BMP receptor proteins, or active fragments thereof, such as CFK1-23a or CFK1-43a. These cells or compositions may be produced by co-expressing each protein in a selected host cell and isolating the cells in a composition or, in the case where truncated receptor proteins are produced, by isolating the truncated receptor proteins from the culture medium.

As a further aspect of this invention a cell line is provided which comprises a first polynucleotide sequence encoding a first receptor protein, truncated receptor protein, or active fragment thereof and a second polynucleotide sequence encoding a second receptor protein, truncated receptor protein, or active fragment thereof, the sequences being under control of one or more suitable expression regulatory systems capable of co-expressing the receptor proteins. The cell line may be transfected with one or more than one polynucleotide molecule. Alternatively, the cell line may be a hybrid cell line created by cell fusion as described above.

Another aspect of the invention is a polynucleotide molecule or plasmid vector comprising a polynucleotide sequence encoding a first selected receptor protein, truncated receptor protein, or active fragment thereof and a polynucleotide sequence encoding a second selected receptor protein, truncated receptor protein, or active fragment thereof. The sequences are under the control of at least one suitable regulatory sequence capable of directing co-expression of each protein or active fragment. The molecule may contain



a single transcription unit containing a copy of both genes, or more than one transcription unit, each containing a copy of a single gene.

One embodiment of the method of the present invention for producing compositions of cells or recombinant receptor proteins involves culturing a suitable cell line, which has been co-transfected with a DNA sequence coding for expression of a first receptor protein, truncated receptor protein, or active fragment thereof and a DNA sequence coding for expression of a second receptor protein, truncated receptor protein, or active fragment thereof, under the control of known regulatory sequences. The transformed host cells are cultured and the cells are isolated and purified to form compositions of transformed cells. In the embodiment wherein truncated receptor proteins are produced, the truncated receptor protein is recovered and purified from the culture medium and can be used to form compositions of truncated receptor protein.

In another embodiment of this method which is the presently preferred method of expression of the recombinant receptor proteins of this invention, a single host cell, e.g., a CHO DUKX cell, is co-transfected with a first DNA molecule containing a DNA sequence encoding one receptor protein, such as the receptor protein CFK1-10a, and a second DNA molecule containing a DNA sequence encoding a second selected receptor protein, such as the BMP receptor protein CFK1-23a or CFK1-43a. One or both plasmids contain a selectable marker that can be used to establish stable cell lines expressing the receptor proteins. These separate plasmids containing distinct receptor genes on separate transcription units are mixed and transfected into the CHO cells using conventional protocols. A ratio of plasmids that gives maximal expression of activity in the binding assay can be determined.

For example, equal ratios of a plasmid containing the first receptor protein gene and a dihydrofolate reductase (DHFR) marker gene and another plasmid containing a second receptor protein gene and a DHFR marker gene can be co-introduced into DHFR-deficient-CHO cells, DUKX-BII, by calcium phosphate coprecipitation and transfection, electroporation, microinjection, protoplast fusion or lipofection. Individual DHFR expressing transformants are selected for growth in alpha media with dialyzed fetal calf serum by conventional means. DHFR+ cells containing increased gene copies can be

selected for propagation in increasing concentrations of methotrexate (MTX) (e.g. sequential steps in 0.02, 0.1, 0.5 and 2.0 uM MTX) according to the procedures of Kaufman and Sharp, J. Mol. Biol., 159:601-629 (1982); and Kaufman et al, Mol. Cell Biol., 5:1750 (1983). Expression of or at least one receptor protein linked to DHFR should increase with increasing levels of MTX resistance. Cells that stably express either or both receptor protein/DHFR genes will survive. However at a high frequency, cell lines stably incorporate and express both plasmids that were present during the initial transfection. The conditioned medium is thereafter harvested and the receptor protein isolated by conventional methods and assayed for activity. This approach can be employed with DHFR-deficient cells.

As an alternative embodiment of this method, a DNA molecule containing one selected receptor gene may be transfected into a stable cell line which already expresses another selected receptor gene. For example, a stable CHO cell line expressing the gene for receptor CFK1-10a with the DHFR marker may be transfected with a plasmid containing the gene for receptor gene for CFK1-23a and a second selectable marker gene, e.g., neomycin resistance (Neo). After transfection, the cell is cultured and suitable cells selected by treatment with MTX and the antibiotic, G-418. Surviving cells are then screened for the expression of both receptor proteins. This expression system has the advantage of permitting a single step selection.

Alternative dual selection strategies using different cell lines or different markers can also be used. For example, the use of an adenosine deaminase (ADA) marker to amplify the second receptor gene in a stable CHO cell line expressing a different receptor with the DHFR marker may be preferable, since the level of expression can be increased using deoxycoformycin (DCF)-mediated gene amplification. Alternatively, any cell line expressing a receptor made by first using this marker can then be the recipient of a second receptor expression vector containing a distinct marker and selected for dual resistance and receptor coexpression.

Still another embodiment of a method of expressing the receptors of this invention includes transfecting the host cell with a single DNA molecule encoding multiple genes for expression either on a single transcription unit or on separate transcription units.



Multicistronic expression involves multiple polypeptides encoded within a single transcript, which can be efficiently translated from vectors utilizing a leader sequence, e.g., from the EMC virus, from poliovirus, or from other conventional sources of leader sequences. Two receptor genes (Rx and Ry, respectively) and a selectable marker can be expressed within a single transcription unit. For example, vectors containing the configuration Rx-EMC-Ry-DHFR or Rx-EMC-Ry-EMC-DHFR can be transfected into CHO cells and selected and amplified using the DHFR marker. A plasmid may be constructed which contains DNA sequences encoding two different receptors, one or more marker genes and a suitable leader or regulatory sequence on a single transcription unit.

Similarly, host cells may be transfected with a single plasmid which contains separate transcription units for each receptor. A selectable marker, e.g., DHFR, can be contained on a another transcription unit, or alternatively as the second cistron on one or both of the receptor genes. These plasmids may be transfected into a selected host cell for expression of the receptors.

Another embodiment of this expression method involves cell fusion. Two stable cell lines which express selected receptors, such as a cell line transformed with a vector for CFK1-23a (e.g., pMV23a) and a cell line stably transformed with a vector for CFK1-43a (e.g., pMV43a), developed using the DHFR/MTX gene amplification system and expressing receptors at high levels, can be transfected with one of several dominant marker genes (e.g., neo<sup>r</sup>, hygromycin<sup>r</sup>, GPT). After sufficient time in coculture (approximately one day) one resultant cell line expressing one receptor and a dominant marker can be fused with a cell line expressing a different receptor and preferably a different marker using a fusogenic reagent, such as polyethylene glycol, Sendai virus or other known agent.

The resulting cell hybrids expressing both dominant markers and DHFR can be selected using the appropriate culture conditions, and screened for coexpression of the receptors, truncated receptors or their fragments. The selected hybrid cell contains sequences encoding both selected receptors, and both receptors will be retained within the membrane of the cell. Compositions of the cells expressing multiple receptors can be used in the methods of the present invention to interact with BMP. In the case where genes encoding truncated receptors are used, the truncated receptor is formed in the cell and then

secreted. The truncated receptor protein is obtained from the conditioned medium and isolated and purified therefrom by conventional methods. The resulting receptor protein composition may be characterized by methods described herein and may be used in the methods of the present invention, for example, to compete with receptors present in cells for ligand binding and thus inhibit the activity of BMP.

Cell lines generated from the approaches described above can be used to produce co-expressed receptor polypeptides. The receptor proteins are retained within the membrane of the cells. Compositions of the cells may be used in order to increase response to BMP, for example to increase cartilage and/or bone formation. Compositions of the cells may be applied in conjunction with BMP.

Where truncated receptor polypeptides are produced, the receptor proteins are isolated from the cell medium in a form substantially free from other proteins with which they are co-produced as well as from other contaminants found in the host cells by conventional purification techniques. The presently preferred method of production is co-transfection of different vectors into CHO cells and methotrexate-mediated gene amplification. Stable cell lines may be used to generate conditioned media containing truncated receptor that can be purified and assayed for *in vitro* and *in vivo* activities. For example, the resulting truncated receptor-producing cell lines obtained by any of the methods described herein may be screened for activity by the binding assays described in Example VI, RNA expression, and protein expression by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE).

The above-described methods of co-expression of the receptors of this invention utilize suitable host cells or cell lines. Suitable cells preferably include mammalian cells, such as Chinese hamster ovary cells (CHO). The selection of suitable mammalian host cells and methods for transformation, culture, amplification, screening and product production and purification are known in the art. See, e.g., Gething and Sambrook, *Nature*, 293:620-625 (1981), or alternatively, Kaufman et al, *Mol. Cell. Biol.*, 5(7):1750-1759 (1985) or Howley et al, U. S. Patent 4,419,446. Other suitable mammalian cell lines are the CV-1 cell line, BHK cell lines and the 293 cell line.

Another aspect of the present invention provides DNA molecules or plasmid vectors

for use in expression of these recombinant receptor proteins. These plasmid vectors may be constructed by resort to known methods and available components known to those of skill in the art. In general, to generate a vector useful in the methods of this invention, the DNA encoding the desired receptor protein, truncated receptor protein, or active  
5 fragment thereof, is transferred into one or more appropriate expression vectors suitable for the selected host cell.

It is presently contemplated that any expression vector suitable for efficient expression in mammalian cells may be employed to produce the recombinant receptor proteins of this invention in mammalian host cells. Preferably the vectors contain the  
10 selected receptor DNA sequences described above and in the Sequence Listings, which encode selected receptor proteins, or truncated receptor proteins. Alternatively, vectors incorporating modified sequences are also embodiments of the present invention and useful in the production of the vectors.

In addition to the specific vectors described above, one skilled in the art can  
15 construct mammalian expression vectors by employing the sequence of SEQ ID NO:1,3,5,7, or 9 or other DNA sequences coding for receptor proteins, truncated receptor proteins, or active fragments thereof and known vectors, such as pCD [Okayama et al, Mol. Cell Biol., 2:161-170 (1982)] and pJL3, pJL4 [Gough et al, EMBO J., 4:645-653 (1985)]. The receptor DNA sequences can be modified by removing the non-coding  
20 nucleotides on the 5' and 3' ends of the coding region. The deleted non-coding nucleotides may or may not be replaced by other sequences known to be beneficial for expression. The transformation of these vectors into appropriate host cells as described above can produce desired receptor proteins.

One skilled in the art could manipulate the sequences of SEQ ID NO:1,3,5,7, or  
25 9 by eliminating or replacing the mammalian regulatory sequences flanking the coding sequence with e.g., yeast or insect regulatory sequences, to create vectors for intracellular or extracellular expression by yeast or insect cells. [See, e.g., procedures described in published European Patent Application 155,476] for expression in insect cells; and procedures described in published PCT application WO86/00639 and European Patent  
30 Application EPA 123,289 for expression in yeast cells].

Similarly, bacterial sequences and preference codons may replace sequences in the described and exemplified mammalian vectors to create suitable expression systems for use in the production of receptor proteins in the method described above. For example, the coding sequences could be further manipulated (e.g., ligated to other known linkers or modified by deleting non-coding sequences therefrom or altering nucleotides therein by other known techniques). The modified receptor coding sequences could then be inserted into a known bacterial vector using procedures such as described in T. Taniguchi et al, Proc. Natl. Acad. Sci. USA, 77:5230-5233 (1980). The exemplary bacterial vector could then be transformed into bacterial host cells and receptor proteins expressed thereby.

Other vectors useful in the methods of this invention may contain multiple genes in a single transcription unit. For example, a proposed plasmid contains the CFK1-10a receptor gene followed by the EMC leader sequence, followed by the CFK1-23a BMP receptor gene, followed by the DHFR marker gene. Another example contains the CFK1-23a BMP receptor gene, the EMC leader, the W101 serine/threonine kinase receptor gene, another EMC leader sequence and the DHFR marker gene. Alternatively, the vector may contain more than one transcription unit. As one example, the plasmid may contain a transcription unit for CFK1-23a BMP receptor gene and a separate transcription unit for CFK1-43a receptor gene, i.e., CFK1-23a-EMC-DHFR and CFK1-43a-EMC-DHFR. Alternatively, each transcription unit on the plasmid may contain a different marker gene. For example, the plasmid may contain CFK1-10a-EMC-Neo and CFK1-43a-EMC-DHFR. Of course, the above examples are not limiting. Other combinations (i.e., co-expression) of the receptors of the present invention are also within the invention.

Additionally the vectors also contain appropriate expression control sequences which are capable of directing the replication and expression of the receptor in the selected host cells. Useful regulatory sequences for such vectors are known to one of skill in the art and may be selected depending upon the selected host cells. Such selection is routine and does not form part of the present invention. Similarly, the vectors may contain one or more selection markers, such as the antibiotic resistance gene, Neo or selectable markers such as DHFR and ADA. The presently preferred marker gene is DHFR. These marker genes may also be selected by one of skill in the art.



The following examples illustrate practice of the present invention in recovering and characterizing the receptor proteins of the present invention and employing them to recover the corresponding human receptor proteins of the present invention, and in expressing the proteins via recombinant techniques.

5     Antibodies

As used herein, a molecule is said to be "BMP-like" if it exhibits at least one BMP-like activity. For the purposes of the invention, BMP-like activity includes the ability to bind to a truncated BMP receptor and to stimulate growth of BMP-dependent cell lines such as the W-20-17 cell line described in Example VIII. As used herein, the term "BMP-like monoclonal antibody" includes non-human BMP-like monoclonal antibodies, complementarity determining regions (CDRs) of the non-human BMP-like monoclonal antibodies, BMP recognition sites of the non-human BMP-like monoclonal antibodies, all engrafted forms of the BMP recognition sites of the non-human BMP-like monoclonal antibodies, and small peptide BMP analogues.

15     In accordance with the present invention, the term "BMP analogue" encompasses monoclonals, small peptides, and small molecules which possess BMP-like activity. For the purposes of the invention, BMP-like activity is defined as the ability to bind to the truncated BMP receptor and to stimulate growth of BMP-dependent cell lines such as the W-20-17 cell line described in Example VIII.

20     The BMP-like monoclonal antibodies of the invention may be molecularly altered to enhance their utility as human pharmaceuticals. For example, the CDRs of the BMP-like monoclonals comprise specific BMP receptor recognition sites, which are encompassed in the present invention. These CDRs may be engrafted onto human immunoglobulin framework regions to minimize antigenicity caused by the presence of  
25     non-human immunoglobulin regions, for example by the techniques disclosed in WO 91/09967. The BMP receptor recognition sites of the present invention may also be engrafted onto other protein frameworks to maximize the therapeutic efficiency of the BMP analogues developed therefrom. The CDRs of the BMP-like monoclonals may also be molecularly altered to form single chain antibodies.

30     The CDRs of the BMP-like monoclonal antibodies may also be used to make small

peptide BMP analogues. The entire CDR may be present in the small peptide BMP analogues of the invention, or an BMP-like portion of the CDR may comprise such small peptide BMP analogues. Two or more CDRs may be joined in "head-to-head", "head-to-tail", or "tail-to-tail" orientation to form dimer or multimer small peptide BMP analogues.

5 Since each BMP-like monoclonal antibody may contain multiple CDRs, a single CDR may be present in the small peptide BMP analogue or different BMP-like CDRs from one or ~~more BMP-like monoclonal antibodies may be present.~~ Non-naturally occurring, synthetic, and D-amino acids may be substituted for specific amino acids of the BMP-like CDRs.

10 The invention further encompasses peptides which specifically bind the truncated receptor and have BMP-like activity. These peptides may be, but are not exclusively, based on the sequence of the CDRs of BMP-like antibodies. The peptides may be "bridged" or "joined" to other peptides in multimeric structures of two peptides or more to elicit the BMP-like activity. The amino acids of the peptides may be, but are not  
15 exclusively, "unnatural" amino acids of, e.g., D-stereo-specificity or analogues of amino acids with other chemical groups attached to the peptide backbone. The peptides may also be cyclic in structure. A peptide as used herein is a molecule comprising of at least two amino acids and up to thirty amino acids.

The invention further encompasses small organic molecules, which may include  
20 amino acid-like molecules, which exhibit specific binding to the truncated human BMP receptor and which possess BMP-like activity. "Small organic molecules" are defined in accordance with the present invention as non-protein carbon-containing molecules of molecular weight less than 3000 which have been first identified for use as BMP analogues using the truncated receptor of the invention. The small organic molecules of the  
25 invention are capable of being incorporated into oral pharmaceuticals for treatment of bone and/or cartilage disorders.

Pharmaceutical compositions containing the BMP-like monoclonal antibodies, small  
30 ~~peptide BMP analogues, or small organic molecules of the present invention~~ are useful in treating a variety of bone and/or cartilage disorders of diverse etiologies. Such pharmaceutical compositions may also contain pharmaceutically acceptable carriers,



diluents, fillers, salts, buffers, stabilizers, and/or other materials well known in the art. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredient(s). The characteristics of the carrier or other material will depend on the route of administration.

5 Administration of the BMP-like monoclonal antibodies, small peptide BMP analogues or small organic molecules of the invention can be carried out in a variety of conventional ways, including via matrices and/or carriers such as are disclosed in U.S. Patent 5,171,579. For the BMP-like monoclonal antibodies, intravenous administration to the patient is preferred. Cutaneous or subcutaneous injection may also be employed for  
10 administration of the monoclonal antibody embodiment of the invention. Oral administration is preferred for administration of the small peptide and small organic molecule BMP analogue embodiments of the invention.

The amount of BMP-like monoclonal antibody, small peptide analogue or small organic molecule in the pharmaceutical composition of the present invention will depend  
15 upon the nature and severity of the condition being treated, and on the nature of prior treatments which the patient has undergone. Ultimately, the attending physician will decide the amount of BMP analogue with which to treat each individual patient. It is contemplated that the various pharmaceutical compositions of the present invention should contain about 0.1  $\mu$ g to about 100  $\mu$ g of BMP analogue molecule per kg body weight.

20 **EXAMPLES**

**Example I. Identification of the murine KDA-B5 sequence.**

Two peptide sequences were derived from the amino acid sequence of the activin receptor in the region described as the kinase domain of the molecule (Mathews et al., *Cell*, 65:973-982(1991)). These particular sequences were selected on the basis of a  
25 comparison between the amino acid sequence of the Daf-1 gene product and the activin receptor and are predicted to be conserved in other serine threonine kinase receptor molecules. The two peptide sequences selected were:

1. Asn-Glu-Tyr-Val-Ala-Val-Lys
2. His-Arg-Asp-Ile-Lys-Ser

30 The following oligonucleotide primer was designed on the basis of amino acid

sequence #1 and synthesized on an automated DNA synthesizer.

Oligonucleotide primer A: GCGGATCCGARTAYGTNGCNGTNAAR

The first 8 nucleotides of this primer (underlined) comprise a recognition sequence for the restriction endonuclease BamHI in order to facilitate subsequent manipulations of amplified DNA products and are not derived from amino acid sequence #1.

The following oligonucleotide primers were designed on the basis of amino acid sequence #2 and synthesized on an automated DNA synthesizer.

Oligonucleotide primer B: GACTCTAGARCTYTTDATRTCYCTRTG

Oligonucleotide primer C: GACTCTAGARCTYTTDATRTCNCGRTG

Oligonucleotide primer D: GACTCTAGANGAYTTDATRTCYCTRTG

Oligonucleotide primer E: GACTCTAGANGAYTTDATRTCNCGRTG

The first 9 nucleotides of primers B through E (underlined) comprise a recognition sequence for the restriction endonuclease XbaI in order to facilitate subsequent manipulations of amplified DNA products and are not derived from amino acid sequence #2.

The standard nucleotide symbols in the above identified primers are as follows: A=adenosine, C=cytosine, G=guanine, T=thymine, R=adenosine or guanine, Y=cytosine or thymine and D=guanine, adenosine or thymine.

These oligonucleotides have been selected for their predicted ability to specifically amplify serine/threonine kinase domain encoding sequences similar to those found in the activin receptor sequence. Since activin and the BMP molecules are members of the large TGF- $\beta$  superfamily of growth and differentiation factors we predict that their corresponding receptors may also be related to each other in structure and primary amino acid sequence. The TGF- $\beta$  type II receptor sequence (Lin et al., *Cell*, 68:775-785 (1992)) indicates that like the activin receptor it is also a serine threonine kinase. On the basis of the above described relationships, we predicted that these degenerate oligonucleotides will specifically amplify sequences encoding fragments of other serine/threonine kinase receptor molecules including activin receptors, TGF- $\beta$  receptors and BMP receptors.

The BMP-2 responsive mouse cell line W-20-17 was selected as a source of mRNA which we would predict to contain molecules that are capable of encoding BMP receptors.

Total RNA was extracted from W-20-17 cells using established procedures known to those skilled in the art, and mRNA was subsequently selected for by oligo (dT) cellulose chromatography. 10 ng of the W-20-17 mRNA was utilized as a template to synthesize first strand cDNA in a reaction mixture (20  $\mu$ l total volume) containing 1 mM each deoxynucleotide triphosphate (dATP, dGTP, dCTP, dTTP), 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 5 mM MgCl<sub>2</sub>, 1 U/ $\mu$ l RNase inhibitor, 2.5 U/ $\mu$ l reverse transcriptase, 2.5  $\mu$ M random hexmers and 20 ng of the W-20-17 mRNA described above. This reaction mixture was incubated for 10 minutes at room temperature, followed by 15 minutes at 42°C and then 5 minutes at 99°C. The completed first strand cDNA reaction was then placed at 4°C.

Oligonucleotide combinations consisting of oligonucleotide primer A paired with either oligonucleotide primer B, C, D or E were utilized as primers to allow the amplification of specific nucleotide sequences from the first strand W-20-17 cDNA template described above. The amplification reaction was performed by adjusting the first strand cDNA reaction described above (20  $\mu$ l) to a volume of 100  $\mu$ l in order to bring the components of the reaction buffer to the following final concentrations: 2 mM MgCl<sub>2</sub>, 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 2.5 U/100  $\mu$ l Taq DNA polymerase, 1 pM/ $\mu$ l oligonucleotide primer A and 1 pM/ $\mu$ l of either oligonucleotide primer B, C, D or E. The entire reaction mixture was then incubated at 95°C for two minutes and then subjected to thermal cycling in the following manner: 1 minute at 95°C, 1 minute at 40°C and 1 minute at 72°C for forty cycles; followed by a 7 minute incubation at 72°C, after which the completed reaction is held at 4°C.

The DNA which was specifically amplified by this reaction was ethanol precipitated, digested with the restriction endonucleases BamHI and XbaI and subjected to agarose gel electrophoresis. Regions of the gel in which DNA bands were evident were excised and the DNA contained within was eluted with a QIAEX Gel Extraction Kit (Qiagen catalog no: 20020) according to the instructions supplied by the manufacturer. The gel-extracted DNA fragments were subcloned into the plasmid vector pGEM-3 between the BamHI and XbaI sites of the polylinker. DNA sequence analysis of one of the resulting subclones named KDA-B5 indicated that the specifically amplified DNA

sequence insert encodes an amino acid sequence homologous to the corresponding region of the activin receptor and Daf-1 gene product kinase domains in the region between where the oligonucleotide primers A, B, C, D and E were designed (as described in the beginning of this section). The amino acid sequence encoded by this region of the specifically amplified KDA-B5 sequence is 41% and 47% identical to the corresponding regions of the activin receptor and Daf-1 kinase domains, respectively.

—The DNA sequence and derived amino acid sequence of the specifically amplified KDA-B5 DNA fragment are set forth in SEQ ID NO:11 and SEQ ID NO:12, respectively.

Nucleotides 1-24 of the sequence set forth in SEQ ID NO:11 comprise a portion of the oligonucleotide primer A and nucleotides 319-341 comprise a portion of the reverse complement of oligonucleotide primer B utilized to perform the specific amplification reaction. Due to the function of oligonucleotides A and B in initiating the amplification reaction, they may not correspond exactly to the actual sequence encoding the mouse KDA-B5 protein and are therefore not translated in the above amino acid derivation.

#### Example II. Isolation of W101 and W120 clones from a W-20-17 cDNA library.

The 341 bp sequence of the KDA-B5 insert set forth in SEQ ID NO:11 was utilized as a probe to screen a W-20-17 cDNA library under reduced stringency conditions (4X SSC, 0.1% SDS at 60°C) in an attempt to isolate other mouse sequences related to KDA-B5 in the following manner.

1,000,000 recombinants of a W-20-17 (Thies et al., *Endocrinology*, 130:1318-1324 (1992)) cDNA library constructed in the vector  $\lambda$ ZAPII were plated at a density of 20,000 recombinant bacteriophage plaques per plate on 100 plates. Duplicate nitrocellulose replicas of the plates were made. A DNA fragment corresponding to the 341 bp sequence of the KDA-B5 insert set forth in SEQ ID NO:11 was <sup>32</sup>P-labelled by the random priming procedure of Feinberg et al., *Anal. Biochem.* 132:6-13 (1983), and hybridized to one set of filters in standard hybridization buffer (SHB: 5X SSC, 0.1% SDS, 5X Denhardt's, 100  $\mu$ g/ml salmon sperm DNA) at 60°C for 2 days. The other set of filters was hybridized to a DNA probe corresponding to nucleotides # 710 to 1044 of the published sequence of the activin receptor (Mathews et al., *Cell*, 65:973-982 (1991)) under the same conditions described for the first set. This region of the activin receptor kinase domain corresponds



to the DNA sequence of the KDA-B5 insert. The filters were washed under reduced stringency conditions (4X SSC, 0.1% SDS at 60°C). 13 positively hybridizing recombinant bacteriophage plaques were selected and replated for secondaries. Duplicate nitrocellulose replicas of the recombinant plaques from these plates were made. Again, one set of filters was hybridized to the KDA-B5 probe and the other set to the activin receptor probe as described above in the primary screen (in SHB at 60°C for 2 days). Both sets of filters were washed under the reduced stringency conditions described above (4X SSC, 0.1% SDS at 60°C). Two recombinants which hybridized strongly to the KDA-B5 probe but not to the corresponding activin receptor probe were selected for further analysis. These two cDNA clones, designated W-101 and W-120, were plaque purified and their inserts were transferred to the plasmid Bluescript SK (+/-) according to the in vivo excision protocol described by the manufacturer (Stratagene). DNA sequence analysis of these recombinants indicated that they encode proteins homologous to the activin receptor and the Daf-1 gene product. The DNA sequence and derived amino acid sequence of a portion of pMT101 (ATCC 69379) is set forth in SEQ ID NO:7 and SEQ ID NO:8, respectively.

The nucleotide sequence of clone W-101 indicates that it encodes a partial polypeptide of 467 amino acids comprising the carboxy-terminal portion of the murine receptor molecule W-101. A random primed cDNA library was made from W-20-17 mRNA using a random hexmer pd(N)<sub>6</sub> (Pharmacia/LKB catalog # 27-2166-01) to prime synthesis of first strand cDNA from the W-20-17 mRNA template. The library was screened with a 30 base oligonucleotide corresponding to nucleotide # 245 to 274 of SEQ ID NO:7. The DNA sequence utilized to design this oligonucleotide probe was derived from the coding sequence of clone W-101. Hybridization was performed in SHB at 65°C and stringent wash conditions of 0.2X SSC, 0.1% SDS AT 65°C were employed to remove non-specifically bound probe. The DNA insert of one of these clones, WR9, was characterized by DNA sequence analysis and determined to contain additional 5' coding sequences of the murine W-101 receptor not present in the original cDNA clone described above. The 0.7 kb insert of the W9 clone however lacks sequences which encode the C-terminal region of the murine W-101 receptor protein. In order to construct a cDNA



sequence which would encode the complete W-101 receptor protein, DNA sequences from both the original oligo (dT) and the random primed clone W9 have been joined at a common BstEII restriction endonuclease site. The sequence set forth in SEQ ID NO:7 contains an open reading frame of 1515 base pairs which encodes the complete 505 amino acid murine W-101 receptor protein. Nucleotides 1-660 of the sequence set forth in SEQ ID NO:7 are derived from the W9 cDNA clone while the remaining sequence (nucleotides 661-1648) are derived from the oligo (dT) primed cDNA clone W-101. The BstEII restriction endonuclease recognition sequence GGTNACC (located at position 660-666) facilitated the construction of this chimeric cDNA sequence. This construction was accomplished by digesting clone W-101 and clone W9 with the restriction endonuclease BstEII resulting in the linearization of each plasmid at the common BstEII site of their respective inserts. The two linearized plasmids were gel isolated, then ligated together at this site and digested with the restriction endonuclease SalI in order to separate the sequence set forth in SEQ ID NO:7 from the plasmid vector sequences (pBluescript). The DNA fragments resulting from this sequential ligation and SalI digest were electrophoresed on an agarose gel. A region containing a DNA fragment of approximately 2.3 kb, comprising 660 bp of the 5' end of the W9 cDNA and approximately 1.64 kb of the 3' end of the original W-101 cDNA, was excised from the gel and the DNA contained therein was eluted with a QIAEX Gel Extraction Kit (Qiagen catalog no: 20020) according to the instructions supplied by the manufacturer. The resulting DNA fragment which comprises the sequence set forth in SEQ ID NO:7 (encoding the complete murine W101 receptor protein) was subcloned into the mammalian cell expression vector pMT3 at the Sal I site of the polylinker region. This plasmid is designated pMT101.

The nucleotide sequence of a portion of the insert of cDNA clone W120 is set forth in SEQ ID NO:9. The presumed initiator methionine encoding sequence is preceded by 68 bp of 5' untranslated sequence and defines an open reading frame of 1509 bp which encodes the complete 503 amino acid murine receptor molecule W-120. The stop codon is followed by at least 203 bp of 3' untranslated sequence. The insert of clone W120 comprising the sequence set forth in SEQ ID NO:9 is excised from the pBluescript plasmid with the restriction endonuclease EcoRI and transferred to the mammalian cell expression

vector pMT3 at the EcoRI site of the polylinker region. This plasmid is designated pMT120E and has been deposited with the American Type Culture Collection (ATCC #69377).

5 **Example III. Isolation of CFK1-10a, CFK1-23a and CFK1-43a from a CFK1 cDNA library.**

Another BMP-2 responsive cell line CFK1 [Bernier and Goltzman, *J. Cell. Physiol.*, 152:317 (1992)] was selected as a source of mRNA which would be predicted to contain molecules capable of encoding BMP receptors and additional serine/threonine kinase encoding sequences related to the TGF- $\beta$  receptor, the activin receptor, *daf-1*, W-101 and W-120.

1 x 10<sup>6</sup> recombinants of a CFK1 cDNA library constructed in the vector  $\lambda$ ZAPII were plated at a density of 20,000 recombinant bacteriophage plaques per plate on 50 plates. Duplicate nitrocellulose replicas of the plates were made. A 645 base pair DNA fragment of the W-101 cDNA insert corresponding to nucleotides #828-#1472 of SEQ ID NO:7 was <sup>32</sup>P-labelled by the random priming procedure of Feinberg et al., *Anal. Biochem.*, 132:6-13 (1983) and hybridized to both sets of filters in SHB at 60°C for two days. The filters were washed under reduced stringency conditions (4X SSC, 0.1% SDS at 60°C). Many duplicate hybridizing recombinants of various intensities (approximately 200) were noted. 27 bacteriophage plaques which were representative of the broad range of hybridization intensity were plaque purified and their inserts were transferred to the plasmid pBluescript SK (+/-) according to the in vivo excision protocol described by the manufacturer (Stratagene). DNA sequence analysis of several recombinants indicated that they encode proteins homologous to the activin receptor, *Daf-1* and other receptor proteins of the serine/threonine kinase receptor family.

25 The nucleotide sequence of clone CFK1-10a comprises an open reading frame of 1527 bp, encoding a CFK1-10a receptor protein of 509 amino acids. The encoded 509 amino acid CFK1-10a receptor protein is contemplated to be the primary translation product, as the coding sequence is preceded by 458 bp of 5' untranslated sequence with stop codons in all three reading frames. The DNA and derived amino acid sequence of the majority of the insert of CFK1-10a (ATCC # 69380) is set forth in SEQ ID NO:5.

Based on the knowledge of other serine/threonine kinase receptor proteins, the encoded 509 amino acid CFK1-10a has the characteristic features of serine/threonine kinase receptors, particularly those capable of recognizing ligands of the TGF- $\beta$ /BMP superfamily of growth and differentiation factors. This molecule encodes a full length receptor molecule with a characteristic hydrophobic leader sequence which targets the ligand binding domain of the protein to the extracellular space, a transmembrane region which anchors the complete receptor molecule in the cell membrane allowing the positioning of the serine threonine kinase domain within the intracellular space. The ligand binding domain of the CFK1-10a receptor molecule of the invention exhibits a pattern of cysteine conservation noted for other receptors capable of recognizing ligands of the TGF- $\beta$ /BMP superfamily of growth and differentiation factors. The region of the CFK1-10 a receptor protein corresponding to the intracellular serine threonine kinase domain exhibits a significant degree of amino acid sequence identity to the corresponding domain of other receptors of this family as follows. TGF- $\beta$  type II receptor (Genbank Accession No. M85079), 35%; activin type II receptor (Genbank Accession No. M65287), 40%; activin type IIB receptor (Genbank Accession No. M84120), 38%; and *Daf-1* (Genbank Accession No. A35103), 39%. The 3.2 kb insert of the CFK1-10a cDNA clone is excised with the restriction endonuclease NotI and transferred to the mammalian cell expression vector pMV2 at the NotI site of the polylinker. This plasmid is designated CFK1-10a/Not-4.

The CFK1-10a receptor of the present invention is homologous to a human serine/threonine kinase receptor mRNA entered into Genbank, accession number L02911, for which no ligand was identified. This receptor protein is also homologous to the reported murine type I TGF- $\beta$  receptor, Ebner et al., *Science*, 260:1344-1348 (1993)(Genbank Accession No. L15436).

The nucleotide sequence of clone CFK1-23a (ATCC # 69378) comprises an open reading frame of 1596 bp, encoding a CFK1-23a receptor protein of 532 amino acids. The encoded 532 amino acid CFK1-23a receptor protein is contemplated to be the primary translation product. The coding sequence is preceded by 60 bp of 5' untranslated sequence. The DNA and derived amino acid sequence of the majority of the insert of

CFK1-23a is set forth in SEQ ID NO:1.

Based on the knowledge of other serine/threonine kinase receptor proteins, the encoded 532 amino acid CFK1-23a has the characteristic features of serine/threonine kinase receptors, particularly those capable of recognizing ligands of the TGF- $\beta$ /BMP superfamily of growth and differentiation factors. The region of the CFK1-23a receptor protein corresponding to the intracellular serine threonine kinase domain exhibits a significant degree of amino acid sequence identity to the corresponding domain of other receptors of this family as follows: TGF- $\beta$  type II receptor (Genbank Accession No. M85079), 35%; activin type II receptor (Genbank Accession No. M65287), 41%; activin type IIB receptor (Genbank Accession No. M84120), 39%; and *Daf-1* (Genbank Accession No. A35103), 39%. The 2.8 kb insert of the CFK1-23a cDNA clone is excised with the restriction endonuclease EcoRI and transferred to the mammalian cell expression vector pMV2 at the EcoRI site of the polylinker. This plasmid is designated pMV23a.

The nucleotide sequence of clone CFK1-43a comprises an open reading frame of 1506 bp, encoding a CFK1-43a receptor protein of 502 amino acids. The encoded 502 amino acid CFK1-43a receptor protein is contemplated to be the primary translation product, as the coding sequence is preceded by 239 bp of 5' untranslated sequence with stop codons in all three reading frames. The DNA and derived amino acid sequence of the insert of CFK1-43a (ATCC # 69381) is set forth in SEQ ID NO:3.

Based on the knowledge of other serine/threonine kinase receptor proteins, the encoded 502 amino acid CFK1-43a has the characteristic features of serine/threonine kinase receptors, particularly those capable of recognizing ligands of the TGF- $\beta$ /BMP superfamily of growth and differentiation factors. The region of the CFK1-43a receptor protein corresponding to the intracellular serine threonine kinase domain exhibits a significant degree of amino acid sequence identity to the corresponding domain of other receptors of this family as follows: TGF- $\beta$  type II receptor (Genbank Accession No. M85079), 35%; activin type II receptor (Genbank Accession No. M65287), 41%; activin type IIB receptor (Genbank Accession No. M84120), 40%; and *Daf-1* (Genbank Accession No. A35103), 38%. The 2.1 kb insert of the CFK1-43a cDNA clone is excised with the restriction endonuclease EcoRI and transferred to the mammalian cell expression vector



pMV2 at the EcoRI site of the polylinker. This plasmid is designated pMV43a.

The CFK1-43a receptor of the invention is homologous to a chicken serine/threonine kinase receptor mRNA, entered into Genbank Accession No. D 13432, for which no ligand has been identified.

5 **Example IV. Screening for Human BMP receptors.**

Mouse and/or rat BMP receptor genes are presumed to be significantly homologous. Therefore, the mouse coding sequences of W-101 or W-120 or portions thereof can be used to screen a human genomic or human cDNA library or as probes to identify a human cell line or tissue which synthesizes the analogous human BMP receptor proteins. In a similar manner, the rat coding sequences of CFK1-10a, CFK1-23a or CFK1-43a or portions thereof can be utilized to screen human libraries or as probes to identify a human cell line or tissue which synthesize the analogous human BMP receptor proteins. A human genomic library may be screened with such probes, and presumptive positively hybridizing recombinant clones isolated and DNA sequence obtained. Evidence that such recombinants encode portions of the corresponding human BMP receptor proteins relies on the murine or rat/human DNA, protein and gene structure homologies.

Once a recombinant bacteriophage or plasmid containing DNA encoding a portion of a human BMP receptor molecule is obtained, the human coding sequence can be used to identify a human cell line or tissue which synthesizes the corresponding human BMP receptor mRNA. Alternatively, the mouse or rat BMP receptor encoding sequence can be utilized as a probe to identify such a human cell line or tissue. Briefly described, RNA is extracted from a selected cell or tissue source and either electrophoresed on a formaldehyde agarose gel and transferred to nitrocellulose, or reacted with formaldehyde and spotted on nitrocellulose directly. The nitrocellulose is then hybridized to a probe derived from a coding sequence from the mouse, rat or human BMP receptor. Alternatively, the mouse or rat BMP receptor coding sequence is used to design oligonucleotide primers which will specifically amplify a portion of the BMP receptor encoding sequence located in the region between the primers utilized to perform the specific amplification reaction. It is contemplated that mouse, rat and human BMP receptor coding sequences would be sufficiently homologous to allow one to specifically



amplify corresponding human BMP receptor encoding sequences from mRNA, cDNA or genomic DNA templates. Once a positive source has been identified by one of these above described methods, mRNA is selected by oligo (dT) cellulose chromatography and cDNA is synthesized and cloned in a suitable vector (ie.  $\lambda$ gt10,  $\lambda$ ZAPII or other vectors known to those skilled in the art) by established techniques. It is also possible to perform the oligonucleotide primer-directed amplification reaction, described above, directly on a pre-established human-cDNA or genomic library which has been cloned into a  $\lambda$  bacteriophage or plasmid vector. In such cases, a library which yields a specifically amplified DNA product encoding a portion of human BMP receptor protein could be screened directly, utilizing the fragment of amplified BMP receptor encoding DNA as a probe.

#### Example V. Expression of BMP receptors.

In order to produce mouse, rat, human or other mammalian BMP receptor proteins, the DNA encoding it is transferred into an appropriate expression vector (as described above for W-101, W-120, CFK1-10a, CFK1-23a and CFK1-43a) and introduced into mammalian cells or other preferred eukaryotic or prokaryotic hosts by conventional genetic engineering techniques. The DNA sequences encoding BMP receptor protein may be inserted into a vector suitable for a particular host cell, as described above. The preferred expression system for recombinant human BMP receptor proteins is contemplated to be stably transformed mammalian cells.

##### A. COS Cell Expression

As one specific example of expressing a serine/threonine kinase receptor protein of the invention, the insert of CFK1-23a (containing the full length BMP receptor cDNA for CFK1-23a) is released from the vector arms by digestion with EcoRI and subcloned into the EcoRI site of the mammalian expression vector, pMV2, a derivative of pMT2, which has been deposited with ATCC under the accession number ATCC 67122, though other derivatives thereof, such as pMT3, may also be suitable. Plasmid DNA from this subclone is transfected into COS cells by the DEAE-dextran procedure [Sompayrac and Danna PNAS 78:7575-7578 (1981); Luthman and Magnusson, Nucl. Acids Res. 11: 1295-1308 (1983)] and the cells are cultured. Serum-free 24 hr. conditioned medium is

collected from the cells starting 40-70 hr. post-transfection.

## **B. CHO Cell Expression**

### **(1) Serine/threonine kinase Receptor Expression in CHO Cells**

In order to achieve high levels of serine/threonine kinase receptor protein expression, each of 5the DNA sequences of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 or SEQ ID NO:9 are inserted into a eucaryotic expression vector, i.e., pMV2 or pMT3, stably introduced into CHO cells and amplified to high copy number by methotrexate selection of DHFR [Kaufman et al., EMBO J. 6:189 (1987)]. The transformed cells are cultured and the expressed receptor proteins remain 10associated with the cell membrane of the transformed cell. Recombinant receptor proteins of the present invention can be dissociated from the transformed cell membrane and then are recovered and purified from other contaminants present.

A serine/threonine kinase receptor protein of the invention is expressed in CHO cells by releasing the insert of cloned CFK1-43a described above with EcoRI. The insert is subcloned into the EcoRI cloning site of the mammalian expression vector, pMV2 described above, though derivatives 15thereof, i.e., pMT3, may also be suitable.

Methods for producing heterologous protein from CHO cells are known in the art and are described above, at pages 16 through 19.

### **Example VI. Binding assays to determine affinity of cloned receptors for different TGF- $\beta$ /BMP superfamily ligands.**

20 A BMP receptor of the invention can be defined by a protein possessing the ability to bind a particular BMP at a greater binding affinity than TGF- $\beta$ , activin, inhibin or other members of the TGF- $\beta$  family of growth and differentiation factors. The BMP receptors of the present invention bind specifically to a particular BMP such that approximately a 100-fold excess of a competitive ligand such as TGF- $\beta$  or activin will not significantly displace the BMP. Specific binding of BMPs to a particular 25BMP receptor of the invention can be demonstrated by transfecting an expression plasmid containing DNA sequences encoding the particular BMP receptor protein of interest into COS cells and allowing for the transient expression of the BMP receptor protein on the cell surface. Individual BMPs, heterodimeric BMPs or other proteins of the TGF- $\beta$  superfamily are bound to BMP receptor expressing COS cells and the cells are analyzed for their ability to bind specifically to a BMP molecule 30or particular set of BMP proteins with greater affinity than to TGF- $\beta$  or other members of the TGF- $\beta$

superfamily. Such binding assays may be performed in the following manner:

COS cells that have been transfected with an expression vector containing the particular BMP receptor coding sequence of interest (e.g., pMT101, pMT120, CFK1-10a/Not-4, pMV23a or pMV43a) are plated on gelatinized 6 well plates and preincubated at 37°C for 60 minutes in binding buffer (128 mM NaCl, 5 mM KCl, 5 mM MgSO<sub>4</sub>, 1.2 mM CaCl<sub>2</sub>, 50 mM HEPES and 5 mg/ml BSA, pH 7.5). The preincubated COS cells are washed and incubated in binding buffer supplemented with 10 mM KCN and 2 mM NaF for 10 minutes prior to the addition of BMP-4 and/or [<sup>125</sup>I]BMP-4. BMP-4 binding is allowed to equilibrate at 37°C for 60 minutes. Following binding, the cells are washed twice with ice-cold binding buffer and solubilized with 1% Triton X-100, 10% glycerol, 25 mM HEPES and 1 mg/ml BSA, pH 7.5, as described by Massague [Meth. Enzymol. 146: 174-195 (1987)]. Radioactivity is then determined in a gamma counter.

**EXAMPLE VII. Production of truncated receptor proteins for production of truncated protein.**

Truncated receptor proteins of the invention preferably comprise the ligand binding domain but not the transmembrane and serine/threonine kinase domains of the receptor proteins. Such truncated receptor proteins can be expressed in mammalian cells in a manner that the truncated receptor proteins will be secreted into the supernatant rather than be expressed on the surface of the host cell. DNA sequences encoding the ligand binding domain of each receptor protein of the invention can be isolated from DNA sequences encoding the transmembrane and serine/threonine kinase domains of each corresponding receptor protein of the invention and inserted into vectors which will allow for the production of truncated receptor proteins in mammalian cells. Alternatively, the DNA sequences encoding the truncated receptor proteins may be isolated and inserted into suitable vectors for expression in bacterial, insect, viral and yeast cells. Such vectors are known to those skilled in the art and are described elsewhere in this application.

In a preferred embodiment, DNA sequences comprising nucleotides # 61 through # 507 of SEQ ID NO:1 encoding amino acids # 1 through 149 of the CFK1-23a receptor protein of the invention (SEQ ID NO:2) can be specifically amplified and the resulting DNA sequence can be inserted into a standard mammalian cell expression vector (i.e.,

pMV2 or pMT3). This specific amplification can be performed in the following manner:

Oligonucleotide primers comprising the nucleotide sequence # 1 through # 20 of SEQ ID NO:1 and a separate primer comprising the complimentary strand of nucleotide sequence # 488 through 507 of SEQ ID NO:1 are synthesized on an automated DNA synthesizer. Additionally these oligonucleotides could be designed to include recognition sequences of restriction endonucleases known by those skilled in the art (ie. BamHI, EcoRI, XbaI etc.) to be useful in the manipulation of amplified DNA products and the facilitation of their insertion into plasmid vectors. Furthermore the primer comprising nucleotides # 488 through 507 could also include a trinucleotide sequence corresponding to a translational stop codon (ie. TAA, TAG or TGA) in place of nucleotides # 583 through 585 of SEQ ID NO:1. The oligonucleotide comprising nucleotides # 488 through 507 would be designed on the basis of the antisense (complementary) strand of this region of SEQ ID NO: 1. and in combination with an oligonucleotide comprising nucleotides # 1 through 20 of the sense (coding) strand of SEQ ID NO: 1 could be used to specifically amplify a DNA fragment comprising nucleotides #1 through # 507 of SEQ ID NO: 1. This DNA fragment will encode a truncated receptor protein of the invention. The DNA fragment encoding the truncated receptor protein of the invention can be produced by specifically amplifying the sequence comprising nucleotides # 1 through # 507 of SEQ ID NO: 1 through the use of a clone encoding the complete receptor of the invention, such as pMV23a, as a template. This specific DNA amplification reaction can be performed as follows: approximately 1 ng of template DNA, such as pMV23a is combined with 100 pM of an oligonucleotide comprising nucleotides #1 through 20 and 100 pM of an oligonucleotide comprising the complementary strand of nucleotides # 488 through 507 in a 100  $\mu$ l reaction mixture consisting of 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 1.5 mM  $MgCl_2$ , 200  $\mu$ M each deoxynucleotide triphosphate and 2.5 units of Taq DNA polymerase. The entire reaction is then incubated at 95° C for two minutes and then subjected to thermal cycling in the following manner: 1 minute at 95° C, 1 minute at 40° C and 1 minute at 72° C for twenty-five to forty cycles; followed by a 7 minute incubation at 72° C, after which the completed reaction is held at 4° C.

The DNA fragment which is specifically amplified by this reaction is ethanol



precipitated, digested with the appropriate restriction endonucleases in the cases where restriction endonuclease recognition sequences have been added to the oligonucleotides utilized to prime the synthesis of the amplified DNA fragment (described above), and subjected to agarose electrophoresis. A region of the gel in which a DNA band of the expected size is evident is excised and subcloned into a plasmid vector. Alternatively this specifically amplified DNA fragment encoding a truncated receptor protein of the invention could be subclone directly into a standard mammalian cell expression vector such as pMT3 or other such vectors known to those skilled in the art.

Similar manipulations could be performed as above, in order to isolate and express other truncated receptor proteins of the invention.

For example, DNA sequences comprising nucleotides #247 through 618 of SEQ ID NO:3 encoding amino acids # 1 through 124 of the CFK1-43a BMP receptor of the invention (SEQ ID NO:4) can be specifically amplified to produce a DNA fragment comprising nucleotides # 247 through 618 of SEQ ID NO:3 which will encode another truncated BMP receptor protein. These sequences will encode the ligand binding domain of the CFK1-43a BMP receptor but will not encode the transmembrane and serine/threonine kinase domains of the corresponding receptor protein. When inserted into an appropriate expression vector and transfected into the appropriate host cell, this construct will allow the production of a truncated BMP receptor protein which will be secreted into the medium rather than be expressed on the surface of the host cell. This specific amplification reaction can be performed in a similar manner to that described above with respect to the truncated CFK1-23a BMP receptor protein. In this case, oligonucleotides comprising the nucleotide sequence #247 through 266 of SEQ ID NO:3 and a separate oligonucleotide primer comprising the complementary strand of nucleotide sequence # 599 through 618 of SEQ ID NO:3 are utilized to specifically amplify a DNA fragment comprising nucleotides # 247 through 618 of SEQ ID NO:3. These oligonucleotides and a template DNA encoding the corresponding BMP receptor protein of the invention such as pMV43a, are substituted in the specific DNA amplification reaction mixture described earlier.

Additionally, other serine/threonine kinase receptors of the invention such as W-



101, W-120 or CFK1-10a can be produced in a truncated form. the truncated forms of these receptor molecules can be expressed in mammalian cells in a manner that the corresponding truncated proteins will be secreted into the culture media rather than be expressed on the surface of the host cell. The expression of these soluble receptor proteins  
5 can be accomplished through the amplification of DNA fragments encoding the ligand binding domain, but not the transmembrane and serine/threonine kinase domains, of each respective serine/threonine-kinase receptor molecule as described above for the truncated BMP receptor proteins.

#### Example VIII. W-20-17 Bioassays

##### 10 A. Description of W-20-17 cells

Use of the W-20-17 bone marrow stromal cells as an indicator cell line is based upon the conversion of these cells to osteoblast-like cells after treatment with a BMP protein [Thies et al, Journal of Bone and Mineral Research, 5:305 (1990); and Thies et al, Endocrinology, 130:1318 (1992)]. Specifically, W-20-17 cells are a clonal bone  
15 marrow stromal cell line derived from adult mice by researchers in the laboratory of Dr. D. Nathan, Children's Hospital, Boston, MA. Treatment of W-20-17 cells with certain BMP proteins results in (1) increased alkaline phosphatase production, (2) induction of PTH stimulated cAMP, and (3) induction of osteocalcin synthesis by the cells. While (1) and (2) represent characteristics associated with the osteoblast phenotype, the ability to  
20 synthesize osteocalcin is a phenotypic property only displayed by mature osteoblasts. Furthermore, to date we have observed conversion of W-20-17 stromal cells to osteoblast-like cells only upon treatment with BMPs. In this manner, the in vitro activities displayed by BMP treated W-20-17 cells correlate with the in vivo bone forming activity known for BMPs.

25 Below two in vitro assays useful in comparison of BMP activities of formulations of BMPs with the activity of known BMPs are described.

##### B. W-20-17 Alkaline Phosphatase Assay Protocol

W-20-17 cells are plated into 96 well tissue culture plates at a density of 10,000 cells per well in 200  $\mu$ l of media (DME with 10% heat inactivated fetal calf serum, 2 mM  
30 glutamine and 100 Units/ml penicillin + 100  $\mu$ g/ml streptomycin. The cells are allowed

to attach overnight in a 95% air, 5% CO<sub>2</sub> incubator at 37°C.

5       The 200  $\mu$ l of media is removed from each well with a multichannel pipettor and replaced with an equal volume of test sample delivered in DME with 10% heat inactivated fetal calf serum, 2 mM glutamine and 1% penicillin-streptomycin. Test substances are assayed in triplicate.

The test samples and standards are allowed a 24 hour incubation period with the W-20-17 indicator-cells. After the 24-hours, plates are removed from the 37°C incubator and the test media are removed from the cells.

10       The W-20-17 cell layers are washed 3 times with 200  $\mu$ l per well of calcium/magnesium free phosphate buffered saline and these washes are discarded.

15       50  $\mu$ l of glass distilled water is added to each well and the assay plates are then placed on a dry ice/ethanol bath for quick freezing. Once frozen, the assay plates are removed from the dry ice/ethanol bath and thawed at 37°C. This step is repeated 2 more times for a total of 3 freeze-thaw procedures. Once complete, the membrane bound alkaline phosphatase is available for measurement.

50  $\mu$ l of assay mix (50 mM glycine, 0.05% Triton X-100, 4 mM MgCl<sub>2</sub>, 5 mM p-nitrophenol phosphate, pH = 10.3) is added to each assay well and the assay plates are then incubated for 30 minutes at 37°C in a shaking waterbath at 60 oscillations per minute.

20       At the end of the 30 minute incubation, the reaction is stopped by adding 100  $\mu$ l of 0.2 N NaOH to each well and placing the assay plates on ice.

The spectrophotometric absorbance for each well is read at a wavelength of 405 nanometers. These values are then compared to known standards to give an estimate of the alkaline phosphatase activity in each sample. For example, using known amounts of p-nitrophenol phosphate, absorbance values are generated. This is shown in Table 2.

25

#### Table 2

Absorbance values for known amounts of BMPs can be determined and converted to  $\mu$ moles of p-nitrophenol phosphate cleaved per unit time as shown in Table 3.

30

#### Table 3

Absorbance Values for Known Standards  
of P-Nitrophenol Phosphate

<u>P-nitrophenol phosphate umoles</u>	<u>Mean absorbance (405 nm)</u>
0.000	0
0.006	0.261 +/- .024
0.012	0.521 +/- .031
0.018	0.797 +/- .063
0.024	1.074 +/- .061
0.030	1.305 +/- .083

Alkaline Phosphatase Values for W-20 Cells  
Treating with BMP-2

<u>BMP-2 concentration</u> <u>ng/ml</u>	<u>Absorbance Reading</u> <u>405 nmeters</u>	<u>umoles substrate</u> <u>per hour</u>
0	0.645	0.024
1.56	0.696	0.026
3.12	0.765	0.029
6.25	0.923	0.036
12.50	1.121	0.044
25.0	1.457	0.058
50.0	1.662	0.067
100.0	1.977	0.080

These values are then used to compare the activities of known amounts of new BMP formulations to known active BMP formulations.

5

C. Osteocalcin RIA Protocol

W-20-17 cells are plated at  $10^6$  cells per well in 24 well multiwell tissue culture

dishes in 2 mls of DME containing 10% heat inactivated fetal calf serum, 2 mM glutamine. The cells are allowed to attach overnight in an atmosphere of 95% air 5% CO<sub>2</sub> at 37°C.

5 The next day the medium is changed to DME containing 10% fetal calf serum, 2 mM glutamine and the test substance in a total volume of 2 ml. Each test substance is administered to triplicate wells. The test substances are incubated with the W-20-17 cells for a total of 96 hours with replacement at 48 hours by the same test medias.

10 At the end of 96 hours, 50 µl of the test media is removed from each well and assayed for osteocalcin production using a radioimmunoassay for mouse osteocalcin. The details of the assay are described in the kit manufactured by Biomedical Technologies Inc., 378 Page Street, Stoughton, MA 02072. Reagents for the assay are found as product numbers BT-431 (mouse osteocalcin standard), BT-432 (Goat anti-mouse Osteocalcin), BT-431R (iodinated mouse osteocalcin), BT-415 (normal goat serum) and BT-414 (donkey anti goat IgG). The RIA for osteocalcin synthesized by W-20-17 cells in response to BMP treatment is carried out as described in the protocol provided by the manufacturer.

#### 15 Example IX. Rat Ectopic Study

Twenty four Long-Evans male rats are divided into 6 test groups. Each receives a subcutaneous implant, 200 uL in size, with either a 0 or 20 ug/100 uL dose of a particular BMP/matrix sample, for example BMP/PLGA porous particles/blood clot, as disclosed in U.S. Patent 5,171,579, the disclosure of which is hereby incorporated by reference. After 14 days, the rats are sacrificed and each animal is evaluated for bone formation.

#### Example X. Polyclonal Antibodies against the Truncated BMP Receptor

25 Three rabbits are injected with purified truncated BMP receptor protein. Polyclonal antibodies from sera of these rabbits are purified by chromatography on a Protein A column. The antibodies are tested for the ability to bind BMP receptors.

#### Example XI. Monoclonal Antibodies Against the Truncated Human BMP Receptor

30 Three sets of three mice are immunized with purified truncated BMP receptor protein. The spleens of the mice are used to generate multiple hybridoma cell lines. The

- conditioned media from non-clonal mouse hybridoma lines are tested for ability to bind to BMP receptors. Clonal lines can be developed from these by sequential serial and limiting dilution cloning. As non-clonal lines become clonal those lines which are positive for activity in the non-clonal stage will retain the ability to produce an antibody which
- 5 binds BMP receptor. Conversely, lines which are negative will stay negative throughout the cloning process. Monoclonal antibodies may be purified from ascites derived from both positive and negative hybridoma clones.



## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: GENETICS INSTITUTE, INC.
- (ii) TITLE OF INVENTION: RECEPTOR PROTEINS
- (iii) NUMBER OF SEQUENCES: 19
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Genetics Institute Inc.- Legal Affairs
  - (B) STREET: 87 CambridgePark Drive
  - (C) CITY: Cambridge
  - (D) STATE: MA
  - (E) COUNTRY: USA
  - (F) ZIP: 02140
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: IBM PC compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
  - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER: TBD
  - (B) FILING DATE: HERewith
  - (C) CLASSIFICATION
- (vii) PRIOR APPLICATION DATA
  - (A) APPLICATION NUMBER: US 08/123,934
  - (B) FILING DATE: 17-SEP-1993
  - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: LAZAR, Steven R
  - (B) REGISTRATION NUMBER: 32,618
  - (C) REFERENCE/DOCKET NUMBER: 5203-PCT
- (ix) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: (617) 498-8260
  - (B) TELEFAX: (617) 876-5851

## (2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1813 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: CFK1-23a
- (ix) FEATURE:
  - (A) NAME/KEY: CDS
  - (B) LOCATION: 61..1656

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CTAGTGGATC	CCCCGGGCTG	CAGGAATTCT	GCGGCCGCCA	GGACACGTGC	GAATTGGACA	60
ATG ACT CAG CTA TAC ACT TAC ATC AGA TTA CTG GGA GCC TGT CTG TTC	108					
Met Thr Gln Leu Tyr Thr Tyr Ile Arg Leu Leu Gly Ala Cys Leu Phe						
1 5 10 15						
ATC ATT TCT CAT GTT CAA GGG CAG AAT CTA GAT AGT ATG CTC CAT GGT	156					
Ile Ile Ser His Val Gln Gly Gln Asn Leu Asp Ser Met Leu His Gly						
20 25 30						
ACT GGT ATG AAA TCA GAC GTG GAC CAG AAG AAG CCG GAA AAT GGA GTG	204					
Thr Gly Met Lys Ser Asp Val Asp Gln Lys Lys Pro Glu Asn Gly Val						
35 40 45						
ACG TTA GCA CCA GAG GAC ACC TTA CCT TTC TTA AAA TGC TAT TGC TCA	252					
Thr Leu Ala Pro Glu Asp Thr Leu Pro Phe Leu Lys Cys Tyr Cys Ser						
50 55 60						
GGA CAC TGC CCA GAT GAC GCT ATT AAT AAC ACA TGC ATA ACT AAT GGC	300					
Gly His Cys Pro Asp Asp Ala Ile Asn Asn Thr Cys Ile Thr Asn Gly						
65 70 75 80						
CAT TGC TTT GCC ATT ATA GAA GAA GAT GAT CAG GGA GAA ACC ACG TTA	348					
His Cys Phe Ala Ile Ile Glu Glu Asp Asp Gln Gly Glu Thr Thr Leu						
85 90 95						
ACT TCT GGG TGT ATG AAG TAT GAA GGC TCT GAT TTT CAA TGC AAG GAT	396					
Thr Ser Gly Cys Met Lys Tyr Glu Gly Ser Asp Phe Gln Cys Lys Asp						
100 105 110						
TCA CCA AAA GCC CAG CTA CGC AGG ACA ATA GAA TGT TGT CCG ACC AAT	444					
Ser Pro Lys Ala Gln Leu Arg Arg Thr Ile Glu Cys Cys Arg Thr Asn						
115 120 125						
TTG TGC AAC CAA TAT TTG CAG CCT ACA CTG CCC CCT GTC GTT ATA GGC	492					
Leu Cys Asn Gln Tyr Leu Gln Pro Thr Leu Pro Pro Val Val Ile Gly						
130 135 140						
CCA TTC TTT GAT GGC AGC GTC CGA TGG CTG GCT GTG CTC ATC TCT ATG	540					
Pro Phe Phe Asp Gly Ser Val Arg Trp Leu Ala Val Leu Ile Ser Met						
145 150 155 160						
GCT GTC TGT ATT GTC GCC ATG ATC GTC TTC TCC AGC TGC TTC TGT TAC	588					
Ala Val Cys Ile Val Ala Met Ile Val Phe Ser Ser Cys Phe Cys Tyr						
165 170 175						
AAA CAT TAC TGT AAG AGT ATC TCA AGC AGA GGT CGT TAC AAC CGT GAC	636					
Lys His Tyr Cys Lys Ser Ile Ser Ser Arg Gly Arg Tyr Asn Arg Asp						
180 185 190						
TTG GAA CAG GAT GAA GCA TTT ATT CCA GTA GGA GAA TCA CTG AAA GAC	684					
Leu Glu Gln Asp Glu Ala Phe Ile Pro Val Gly Glu Ser Leu Lys Asp						
195 200 205						
CTG ATT GAC CAG TCA CAA AGC TCT GGT AGT GGA TCT GGA TTA CCT TTA	732					
Leu Ile Asp Gln Ser Gln Ser Ser Gly Ser Gly Ser Gly Leu Pro Leu						
210 215 220						
TTG GTT CAG CGA ACT ATT GCC AAA CAG ATT CAG ATG GTT CGG CAG GTT	780					
Leu Val Gln Arg Thr Ile Ala Lys Gln Ile Gln Met Val Arg Gln Val						
225 230 235 240						
GGT AAG GGC CGG TAT GGA GAA GTA TGG ATG GGT AAA TGG CGT GGT GAA	828					
Gly Lys Gly Arg Tyr Gly Glu Val Trp Met Gly Lys Trp Arg Gly Glu						
245 250 255						

AAA GTG GCT GTC AAA GTA TTT TTT ACC ACT GAA GAA GCT AGC TGG TTT Lys Val Ala Val Lys Val Phe Phe Thr Thr Glu Glu Ala Ser Trp Phe 260 265 270	876
AGA GAA ACA GAA ATC TAC CAG ACG GTG TTA ATG CGT CAT GAA AAT ATA Arg Glu Thr Glu Ile Tyr Gln Thr Val Leu Met Arg His Glu Asn Ile 275 280 285	924
CTT GGT TTT ATA GCT GCA GAC ATT AAA GGC ACC GGT TCC TGG ACT CAG Leu Gly Phe Ile Ala Ala Asp Ile Lys Gly Thr Gly Ser Trp Thr Gln 290 295 300	972
CTG TAT TTG ATT ACT GAT TAC CAT GAG AAT GGG TCT CTC TAT GAC TTC Leu Tyr Leu Ile Thr Asp Tyr His Glu Asn Gly Ser Leu Tyr Asp Phe 305 310 315 320	1020
CTG AAA TGT GCC ACC CTG GAC ACC AGA GCC CTA CTC AAG TTA GCT TAT Leu Lys Cys Ala Thr Leu Asp Thr Arg Ala Leu Leu Lys Leu Ala Tyr 325 330 335	1068
TCT GCT GCC TGT GGT CTG TGC CAC CTC CAC ACA GAA ATT TAT GGC ACG Ser Ala Ala Cys Gly Leu Cys His Leu His Thr Glu Ile Tyr Gly Thr 340 345 350	1116
CAA GGC AAG CCT GCA ATT GCT CAT CGA GAC CTG AAG AGC AAA AAC ATC Gln Gly Lys Pro Ala Ile Ala His Arg Asp Leu Lys Ser Lys Asn Ile 355 360 365	1164
CTT ATT AAG AAA AAT GGT AGT TGC TGT ATT GCT GAC CTG GGC CTA GCT Leu Ile Lys Lys Asn Gly Ser Cys Cys Ile Ala Asp Leu Gly Leu Ala 370 375 380	1212
GTT AAA TTC AAC AGT GAC ACA AAT GAA GTT GAC ATA CCC TTG AAC ACC Val Lys Phe Asn Ser Asp Thr Asn Glu Val Asp Ile Pro Leu Asn Thr 385 390 395 400	1260
AGG GTG GGC ACC AGG CGG TAC ATG GCT CCA GAA GTG CTG GAC GAG AGC Arg Val Gly Thr Arg Arg Tyr Met Ala Pro Glu Val Leu Asp Glu Ser 405 410 415	1308
CTG AGT AAA AAC CAT TTC CAG CCC TAC ATC ATG GCT GAC ATC TAC AGC Leu Ser Lys Asn His Phe Gln Pro Tyr Ile Met Ala Asp Ile Tyr Ser 420 425 430	1356
TTT GGT TTG ATC ATT TGG GAG ATG GCC CGT CGC TGT ATT ACA GGA GGA Phe Gly Leu Ile Ile Trp Glu Met Ala Arg Arg Cys Ile Thr Gly Gly 435 440 445	1404
ATC GTG GAG GAA TAT CAA TTA CCA TAT TAC AAC ATG GTG CCT AGT GAC Ile Val Glu Glu Tyr Gln Leu Pro Tyr Tyr Asn Met Val Pro Ser Asp 450 455 460	1452
CCA TCT TAT GAA GAC ATG CGT GAG GTC GTG TGT GTG AAA CGC TTG CGG Pro Ser Tyr Glu Asp Met Arg Glu Val Val Cys Val Lys Arg Leu Arg 465 470 475 480	1500
CCA ATC GTC TCT AAC CGC TGG AAC AGT GAT GAA TGT CTT CGA GCC GTT Pro Ile Val Ser Asn Arg Trp Asn Ser Asp Glu Cys Leu Arg Ala Val 485 490 495	1548
TTG AAG CTG ATG TCA GAA TGC TGG GCC CAT AAT CCA GCA TCC AGA CTC Leu Lys Leu Met Ser Glu Cys Trp Ala His Asn Pro Ala Ser Arg Leu 500 505 510	1596
ACA GCT TTG AGA ATC AAG AAG ACG CTC GCA AAG ATG GTT GAA TCC CAG Thr Ala Leu Arg Ile Lys Lys Thr Leu Ala Lys Met Val Glu Ser Gln 515 520 525	1644

GAT GTA AAG ATT TGACAAACAG TTTTGAGAAA GAATTTAGAC TGCAAGAAAT 1696  
 Asp Val Lys Ile  
 530

TCACCCGAGG AAGGGTGGAG TTAGCATGGA CTAGGATGTC GGCTTGGTTT CCAGACTCTC 1756

TCCTCTACCA TCTTCACAGG CTGCTAACAG TAAACCTTTC AGGACTCTGC AGAATGC 1813

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

--- (A)---LENGTH: 532 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Thr Gln Leu Tyr Thr Tyr Ile Arg Leu Leu Gly Ala Cys Leu Phe  
 1 5 10 15

Ile Ile Ser His Val Gln Gly Gln Asn Leu Asp Ser Met Leu His Gly  
 20 25 30

Thr Gly Met Lys Ser Asp Val Asp Gln Lys Lys Pro Glu Asn Gly Val  
 35 40 45

Thr Leu Ala Pro Glu Asp Thr Leu Pro Phe Leu Lys Cys Tyr Cys Ser  
 50 55 60

Gly His Cys Pro Asp Asp Ala Ile Asn Asn Thr Cys Ile Thr Asn Gly  
 65 70 75 80

His Cys Phe Ala Ile Ile Glu Glu Asp Asp Gln Gly Glu Thr Thr Leu  
 85 90 95

Thr Ser Gly Cys Met Lys Tyr Glu Gly Ser Asp Phe Gln Cys Lys Asp  
 100 105 110

Ser Pro Lys Ala Gln Leu Arg Arg Thr Ile Glu Cys Cys Arg Thr Asn  
 115 120 125

Leu Cys Asn Gln Tyr Leu Gln Pro Thr Leu Pro Pro Val Val Ile Gly  
 130 135 140

Pro Phe Phe Asp Gly Ser Val Arg Trp Leu Ala Val Leu Ile Ser Met  
 145 150 155 160

Ala Val Cys Ile Val Ala Met Ile Val Phe Ser Ser Cys Phe Cys Tyr  
 165 170 175

~~Lys His Tyr Cys Lys Ser Ile Ser Ser Arg Gly Arg Tyr Asn Arg Asp~~  
 180 185 190

Leu Glu Gln Asp Glu Ala Phe Ile Pro Val Gly Glu Ser Leu Lys Asp  
 195 200 205

Leu Ile Asp Gln Ser Gln Ser Ser Gly Ser Gly Ser Gly Leu Pro Leu  
 210 215 220

Leu Val Gln Arg Thr Ile Ala Lys Gln Ile Gln Met Val Arg Gln Val  
 225 230 235 240

Gly Lys Gly Arg Tyr Gly Glu Val Trp Met Gly Lys Trp Arg Gly Glu  
 245 250 255

Lys Val Ala Val Lys Val Phe Phe Thr Thr Glu Glu Ala Ser Trp Phe  
 260 265 270  
 Arg Glu Thr Glu Ile Tyr Gln Thr Val Leu Met Arg His Glu Asn Ile  
 275 280 285  
 Leu Gly Phe Ile Ala Ala Asp Ile Lys Gly Thr Gly Ser Trp Thr Gln  
 290 295 300  
 Leu Tyr Leu Ile Thr Asp Tyr His Glu Asn Gly Ser Leu Tyr Asp Phe  
 305 310 315 320  
 Leu Lys Cys Ala Thr Leu Asp Thr Arg Ala Leu Leu Lys Leu Ala Tyr  
 325 330 335  
 Ser Ala Ala Cys Gly Leu Cys His Leu His Thr Glu Ile Tyr Gly Thr  
 340 345 350  
 Gln Gly Lys Pro Ala Ile Ala His Arg Asp Leu Lys Ser Lys Asn Ile  
 355 360 365  
 Leu Ile Lys Lys Asn Gly Ser Cys Cys Ile Ala Asp Leu Gly Leu Ala  
 370 375 380  
 Val Lys Phe Asn Ser Asp Thr Asn Glu Val Asp Ile Pro Leu Asn Thr  
 385 390 395 400  
 Arg Val Gly Thr Arg Arg Tyr Met Ala Pro Glu Val Leu Asp Glu Ser  
 405 410 415  
 Leu Ser Lys Asn His Phe Gln Pro Tyr Ile Met Ala Asp Ile Tyr Ser  
 420 425 430  
 Phe Gly Leu Ile Ile Trp Glu Met Ala Arg Arg Cys Ile Thr Gly Gly  
 435 440 445  
 Ile Val Glu Glu Tyr Gln Leu Pro Tyr Tyr Asn Met Val Pro Ser Asp  
 450 455 460  
 Pro Ser Tyr Glu Asp Met Arg Glu Val Val Cys Val Lys Arg Leu Arg  
 465 470 475 480  
 Pro Ile Val Ser Asn Arg Trp Asn Ser Asp Glu Cys Leu Arg Ala Val  
 485 490 495  
 Leu Lys Leu Met Ser Glu Cys Trp Ala His Asn Pro Ala Ser Arg Leu  
 500 505 510  
 Thr Ala Leu Arg Ile Lys Lys Thr Leu Ala Lys Met Val Glu Ser Gln  
 515 520 525  
 Asp Val Lys Ile  
 530

## (2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 2076 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- (vii) IMMEDIATE SOURCE:
- (B) CLONE: CFK1-43a



## (ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 247..1752

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GCGGCCGCGC CGGCGTGGTG CTCGGAGTGC GGGCGCCGAG GACCCGGGAC CAGGGGCGCG	60
GCGGCGGGTT GGAGTTCAAG GTACTCGTTA CGTGTGACGA GGAAGTGAAG CCCATTCCAT	120
GCCTTGCTGA GAAAGGTTCA AACTTCGGCT GAATCACAAC CATTTGGCGC TGAGCTATGA	180
CAAGAGAGCA AACAAAAAGT TAAAGGAGCA ACTCGGCCAT AAGTGACAGA GAAGTTCGTT	240
GATAAC ATG CTC TTA CGA AGC TCT GGA AAA TTA AAT GTG GGC ACC AAG	288
Met Leu Leu Arg Ser Ser Gly Lys Leu Asn Val Gly Thr Lys	
1 5 10	
AAG GAG GAT GGT GAG AGT ACA GCC CCC ACT GCT CGG CCC AAG GTC CTG	336
Lys Glu Asp Gly Glu Ser Thr Ala Pro Thr Ala Arg Pro Lys Val Leu	
15 20 25 30	
CGT TGT AAA TGC CAC CAC CAC TGT CCT GAA GAC TCA GTC AAC AAT ATC	384
Arg Cys Lys Cys His His His Cys Pro Glu Asp Ser Val Asn Asn Ile	
35 40 45	
TGC AGC ACA GAT GGG TAC TGC TTC ACG ATG ATA GAA GAA GAC GAC TCT	432
Cys Ser Thr Asp Gly Tyr Cys Phe Thr Met Ile Glu Glu Asp Asp Ser	
50 55 60	
GGA ACG CCT GTT GTC ACC TCC GGA TGC CTA GGA CTA GAA GGG TCA GAT	480
Gly Thr Pro Val Val Thr Ser Gly Cys Leu Gly Leu Glu Gly Ser Asp	
65 70 75	
TTT CAA TGT CGC GAC ACG CCC ATC CCT CAT CAG AGA AGG TCA ATT GAA	528
Phe Gln Cys Arg Asp Thr Pro Ile Pro His Gln Arg Arg Ser Ile Glu	
80 85 90	
TGC TGC ACA GAA AGG AAC GAA TGT AAT AAA GAT CTC CAC CCC ACG CTG	576
Cys Cys Thr Glu Arg Asn Glu Cys Asn Lys Asp Leu His Pro Thr Leu	
95 100 105 110	
CCT CCC CTG AAG GAC AGA GAT TTT GTT GAT GGA CCC ATA CAC CAC AAA	624
Pro Pro Leu Lys Asp Arg Asp Phe Val Asp Gly Pro Ile His His Lys	
115 120 125	
GCC TTA CTC ATA TCT GTG ACT GTC TGT AGT TTA CTC TTG GTC CTC ATT	672
Ala Leu Leu Ile Ser Val Thr Val Cys Ser Leu Leu Leu Val Leu Ile	
130 135 140	
ATT TTA TTC TGT TAC TTC AGG TAT AAA AGA CAA GAA GCC AGA CCT CGG	720
Ile Leu Phe Cys Tyr Phe Arg Tyr Lys Arg Gln Glu Ala Arg Pro Arg	
145 150 155	
TAC AGC ATT GGG CTG GAG CAG GAT GAA ACG TAC ATT CCT CCT GGA GAA	768
Tyr Ser Ile Gly Leu Glu Gln Asp Glu Thr Tyr Ile Pro Pro Gly Glu	
160 165 170	
TCC CTG AGA GAC TTG ATT GAG CAA TCG CAG AGC TCG GGA AGT GGC TCA	816
Ser Leu Arg Asp Leu Ile Glu Gln Ser Gln Ser Ser Gly Ser Gly Ser	
175 180 185 190	
GGA CTC CCT CTG CTG GTC CAA AGG ACA ATA GCT AAG CAA ATT CAG ATG	864
Gly Leu Pro Leu Leu Val Gln Arg Thr Ile Ala Lys Gln Ile Gln Met	
195 200 205	

GTG	AAG	CAA	ATT	GGA	AAA	GGT	CGC	TAT	GGC	GAA	GTG	TGG	ATG	GGA	AAG	912
Val	Lys	Gln	Ile	Gly	Lys	Gly	Arg	Tyr	Gly	Glu	Val	Trp	Met	Gly	Lys	
			210					215					220			
TGG	CGT	GGA	GAA	AAG	GTA	GCT	GTG	AAA	GTG	TTC	TTC	ACC	ACG	GAG	GAA	960
Trp	Arg	Gly	Glu	Lys	Val	Ala	Val	Lys	Val	Phe	Phe	Thr	Thr	Glu	Glu	
		225					230					235				
GCC	AGC	TGG	TTC	CGA	GAG	ACT	GAG	ATA	TAT	CAG	ACG	GTC	CTG	ATG	AGG	1008
Ala	Ser	Trp	Phe	Arg	Glu	Thr	Glu	Ile	Tyr	Gln	Thr	Val	Leu	Met	Arg	
	240					245					250					
CAC	GAG	AAC	ATT	CTG	GGG	TTC	ATT	GCA	GCA	GAT	ATC	AAA	GGG	ACT	GGG	1056
His	Glu	Asn	Ile	Leu	Gly	Phe	Ile	Ala	Ala	Asp	Ile	Lys	Gly	Thr	Gly	
	255				260					265					270	
TCT	TGG	ACT	CAG	TTA	TAC	CTC	ATC	ACA	GAC	TAT	CAT	GAA	AAC	GGG	TCT	1104
Ser	Trp	Thr	Gln	Leu	Tyr	Leu	Ile	Thr	Asp	Tyr	His	Glu	Asn	Gly	Ser	
				275					280					285		
CTT	TAT	GAC	TAT	CTG	AAA	TCC	ACC	ACC	TTA	GAT	GCC	AAG	TCC	ATG	CTG	1152
Leu	Tyr	Asp	Tyr	Leu	Lys	Ser	Thr	Thr	Leu	Asp	Ala	Lys	Ser	Met	Leu	
			290					295					300			
AAG	CTA	GCC	TAC	TCG	TCT	GTC	AGC	GGC	CTG	TGC	CAT	CTA	CAC	ACG	GAA	1200
Lys	Leu	Ala	Tyr	Ser	Ser	Val	Ser	Gly	Leu	Cys	His	Leu	His	Thr	Glu	
		305					310					315				
ATC	TTC	AGC	ACT	CAA	GGC	AAG	CCA	GCC	ATT	GCC	CAT	CGG	GAC	TTG	AAA	1248
Ile	Phe	Ser	Thr	Gln	Gly	Lys	Pro	Ala	Ile	Ala	His	Arg	Asp	Leu	Lys	
	320					325					330					
AGT	AAA	AAC	ATC	CTG	GTG	AAG	AAA	AAT	GGA	ACT	TGC	TGC	ATA	GCA	GAC	1296
Ser	Lys	Asn	Ile	Leu	Val	Lys	Lys	Asn	Gly	Thr	Cys	Cys	Ile	Ala	Asp	
	335				340					345					350	
CTG	GGC	CTG	GCT	GTC	AAG	TTC	ATT	AGT	GAC	ACA	AAT	GAG	GTT	GAC	ATT	1344
Leu	Gly	Leu	Ala	Val	Lys	Phe	Ile	Ser	Asp	Thr	Asn	Glu	Val	Asp	Ile	
				355					360					365		
CCA	CCC	AAC	ACC	CGG	GTT	GGC	ACC	AAG	CGC	TAT	ATG	CCT	CCA	GAA	GTG	1392
Pro	Pro	Asn	Thr	Arg	Val	Gly	Thr	Lys	Arg	Tyr	Met	Pro	Pro	Glu	Val	
			370					375					380			
CTG	GAC	GAG	AGC	TTG	AAT	AGA	ACT	CAT	TTC	CAG	TCC	TAC	ATC	ATG	GCT	1440
Leu	Asp	Glu	Ser	Leu	Asn	Arg	Thr	His	Phe	Gln	Ser	Tyr	Ile	Met	Ala	
		385					390					395				
GAC	ATG	TAC	AGC	TTT	GGA	CTC	ATC	CTC	TGG	GAG	ATT	GCA	AGG	AGA	TGT	1488
Asp	Met	Tyr	Ser	Phe	Gly	Leu	Ile	Leu	Trp	Glu	Ile	Ala	Arg	Arg	Cys	
	400					405					410					
GTT	TCT	GGA	GGT	ATA	GTG	GAA	GAA	TAC	CAG	CTT	CCA	TAT	CAC	GAC	CTG	1536
Val	Ser	Gly	Gly	Ile	Val	Glu	Glu	Tyr	Gln	Leu	Pro	Tyr	His	Asp	Leu	
	415				420					425					430	
GTG	CCC	AGT	GAC	CCC	TCT	TAT	GAG	GAC	ATG	AGA	GAA	ATT	GTG	TGT	ATG	1584
Val	Pro	Ser	Asp	Pro	Ser	Tyr	Glu	Asp	Met	Arg	Glu	Ile	Val	Cys	Met	
				435				440					445			
AAG	AAG	TTA	CGG	CCT	TCA	TTC	CCC	AAT	CGA	TGG	AGC	AGT	GAC	GAG	TGC	1632
Lys	Lys	Leu	Arg	Pro	Ser	Phe	Pro	Asn	Arg	Trp	Ser	Ser	Asp	Glu	Cys	
			450					455					460			
CTC	AGG	CAA	ATG	GGG	AAG	CTT	ATG	ACA	GAG	TGC	TGG	GCG	CAT	AAT	CCT	1680
Leu	Arg	Gln	Met	Gly	Lys	Leu	Met	Thr	Glu	Cys	Trp	Ala	His	Asn	Pro	
		465					470					475				

GCC TCC AGG CTG ACG GCC CTG AGA GTT AAG AAA ACA CTT GCC AAA ATG 1728  
 Ala Ser Arg Leu Thr Ala Leu Arg Val Lys Lys Thr Leu Ala Lys Met  
 480 485 490

TCA GAG TCC CAG GAC ATT AAA CTC TGACGTCAGG TACTTGTGGA CAGAGCAAGG 1782  
 Ser Glu Ser Gln Asp Ile Lys Leu  
 495 500

AATTACACAG AAGCATCCTT AGCCCAAGCC TTGAACGTTG ATCTACTGCC CAGTGAGTTC 1842

AGACTTTCCT CTAAGAGAGC AAGCTGGACA GACACAGAGG AACCCAGAAA CACGGCTTCA 1902

CCATGGCTTT CTGAGGAGGG GAAACCATTT GGGTAACTTG TTCAAGATAT GATGCATGTT 1962

GCTTTCTAAG AAAGCCCTGT ATTTTGGGAT TACCATTTTT TTAAAGAAG AAAGATACTT 2022

TAATTTTAC CAAAATAAAA CAAATATTAT AGAAAAAAG CGGCCGCAGA ATTC 2076

## (2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 502 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Leu Leu Arg Ser Ser Gly Lys Leu Asn Val Gly Thr Lys Lys Glu  
 1 5 10 15

Asp Gly Glu Ser Thr Ala Pro Thr Ala Arg Pro Lys Val Leu Arg Cys  
 20 25 30

~~Lys Cys His His His Cys Pro Glu Asp Ser Val Asn Asn Ile Cys Ser~~  
 35 40 45

Thr Asp Gly Tyr Cys Phe Thr Met Ile Glu Glu Asp Asp Ser Gly Thr  
 50 55 60

Pro Val Val Thr Ser Gly Cys Leu Gly Leu Glu Gly Ser Asp Phe Gln  
 65 70 75 80

Cys Arg Asp Thr Pro Ile Pro His Gln Arg Arg Ser Ile Glu Cys Cys  
 85 90 95

Thr Glu Arg Asn Glu Cys Asn Lys Asp Leu His Pro Thr Leu Pro Pro  
 100 105 110

Leu Lys Asp Arg Asp Phe Val Asp Gly Pro Ile His His Lys Ala Leu  
 115 120 125

Leu Ile Ser Val Thr Val Cys Ser Leu Leu Leu Val Leu Ile Ile Leu  
 130 135 140

Phe Cys Tyr Phe Arg Tyr Lys Arg Gln Glu Ala Arg Pro Arg Tyr Ser  
 145 150 155 160

Ile Gly Leu Glu Gln Asp Glu Thr Tyr Ile Pro Pro Gly Glu Ser Leu  
 165 170 175

Arg Asp Leu Ile Glu Gln Ser Gln Ser Ser Gly Ser Gly Ser Gly Leu  
 180 185 190

Pro Leu Leu Val Gln Arg Thr Ile Ala Lys Gln Ile Gln Met Val Lys  
 195 200 205  
 Gln Ile Gly Lys Gly Arg Tyr Gly Glu Val Trp Met Gly Lys Trp Arg  
 210 215 220  
 Gly Glu Lys Val Ala Val Lys Val Phe Phe Thr Thr Glu Glu Ala Ser  
 225 230 235 240  
 Trp Phe Arg Glu Thr Glu Ile Tyr Gln Thr Val Leu Met Arg His Glu  
 245 250 255  
 Asn Ile Leu Gly Phe Ile Ala Ala Asp Ile Lys Gly Thr Gly Ser Trp  
 260 265 270  
 Thr Gln Leu Tyr Leu Ile Thr Asp Tyr His Glu Asn Gly Ser Leu Tyr  
 275 280 285  
 Asp Tyr Leu Lys Ser Thr Thr Leu Asp Ala Lys Ser Met Leu Lys Leu  
 290 295 300  
 Ala Tyr Ser Ser Val Ser Gly Leu Cys His Leu His Thr Glu Ile Phe  
 305 310 315 320  
 Ser Thr Gln Gly Lys Pro Ala Ile Ala His Arg Asp Leu Lys Ser Lys  
 325 330 335  
 Asn Ile Leu Val Lys Lys Asn Gly Thr Cys Cys Ile Ala Asp Leu Gly  
 340 345 350  
 Leu Ala Val Lys Phe Ile Ser Asp Thr Asn Glu Val Asp Ile Pro Pro  
 355 360 365  
 Asn Thr Arg Val Gly Thr Lys Arg Tyr Met Pro Pro Glu Val Leu Asp  
 370 375 380  
 Glu Ser Leu Asn Arg Thr His Phe Gln Ser Tyr Ile Met Ala Asp Met  
 385 390 395 400  
 Tyr Ser Phe Gly Leu Ile Leu Trp Glu Ile Ala Arg Arg Cys Val Ser  
 405 410 415  
 Gly Gly Ile Val Glu Glu Tyr Gln Leu Pro Tyr His Asp Leu Val Pro  
 420 425 430  
 Ser Asp Pro Ser Tyr Glu Asp Met Arg Glu Ile Val Cys Met Lys Lys  
 435 440 445  
 Leu Arg Pro Ser Phe Pro Asn Arg Trp Ser Ser Asp Glu Cys Leu Arg  
 450 455 460  
 Gln Met Gly Lys Leu Met Thr Glu Cys Trp Ala His Asn Pro Ala Ser  
 465 470 475 480  
 Arg Leu Thr Ala Leu Arg Val Lys Lys Thr Leu Ala Lys Met Ser Glu  
 485 490 495  
 Ser Gln Asp Ile Lys Leu  
 500

## (2) INFORMATION FOR SEQ ID NO:5:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3238 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vii) IMMEDIATE SOURCE:

(B) CLONE: CFK1-10a

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 474..2000

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GAATTCTGCG GCCGCGAGGC TGCATTAAGT GGGATATGCC ACCCGTGATT CTGACAGCCG	60
TGACTGCGTG GAGCCTGCTC CGGAACCTCTC CACAGAGGAG CAAAGGAGCT GCCCTCTGTG	120
TCTCCCCGCC CTTCAGCGAG AGTCTGGAAA GAGAACCGAG GTGCTACTGC AGTGGATGAG	180
TAGAGAAGAG TCTGCATCCA GTGCTGGTGA GCTTGTCTGG CTATAGGGAG CCTGCTGGGG	240
GAAACTTACA GCTTCAGAAG ACTCCTGGAG AGCCTCTCCC TCCACACTCT CCCTTTGAGC	300
AGTCAGTGCC TCTCTGCTGG AGAACCTGTG CTGGGTGTGC CCCAGAGCTG GCTTTGACTG	360
TAGCCTGTCA GGCTCTCCCT GGACCTCACG GAACAGCATT GCCAGCCACA CGGCTTCCAA	420
CAAATCACCT CTTTTCATGC TGTTTGGCAC AGATCGAATC TACAGGTTAT ACA ATG	476
	Met 1
GTC GAT GGA GCA ATG ATC CTT TCT GTG CTA ATG ATG ATG GCT CTC CCT	524
Val Asp Gly Ala Met Ile Leu Ser Val Leu Met Met Met Ala Leu Pro	
	5 10 15
TCC CCG AGT ATG GAA GAT GAG GAG CCC AAG GTC AAC CCG AAG CTT TAC	572
<del>Ser-Pro-Ser-Met-Glu-Asp-Glu</del> <del>Glu-Pro-Lys-Val-Asn</del> Pro Lys Leu Tyr	
	20 25 30
ATG TGT GTG TGT GAG GGC CTC TCC TGC GGG AAC GAG GAC CAC TGT GAG	620
Met Cys Val Cys Glu Gly Leu Ser Cys Gly Asn Glu Asp His Cys Glu	
	35 40 45
GGC CAG CAG TGT TTT TCC TCC CTG AGC GTC AAT GAT GGC TTC CGC GTC	668
Gly Gln Gln Cys Phe Ser Ser Leu Ser Val Asn Asp Gly Phe Arg Val	
	50 55 60 65
TAC CAG AAG GGC TGC TTT CAG GTC TAT GAG CAG GGG AAG ATG ACG TGT	716
Tyr Gln Lys Gly Cys Phe Gln Val Tyr Glu Gln Gly Lys Met Thr Cys	
	70 75 80
AAG ACC CCG CCG TCG CCT GGC CAG GCT GTG GAG TGC TGC CAA GGG GAC	764
Lys Thr Pro Pro Ser Pro Gly Gln Ala Val Glu Cys Cys Gln Gly Asp	
	85 90 95
TGG TGC AAC AGG AAC GTC ACG GCC CGG CTG CCC ACT AAA GGG AAA TCC	812
Trp Cys Asn Arg Asn Val Thr Ala Arg Leu Pro Thr Lys Gly Lys Ser	
	100 105 110
TTC CCT GGA TCG CAG AAC TTC CAC CTG GAA GTT GGC CTT ATC ATC CTC	860
Phe Pro Gly Ser Gln Asn Phe His Leu Glu Val Gly Leu Ile Ile Leu	
	115 120 125
TCC GTG GTG TTT GCG GTA TGC CTT TTC GCT TGC ATC CTT GGC GTT GCT	908
Ser Val Val Phe Ala Val Cys Leu Phe Ala Cys Ile Leu Gly Val Ala	
	130 135 140 145



CTC AGG AAG TTT AAA AGG CGC AAT CAA GAG CGC CTG AAC CCC AGA GAC Leu Arg Lys Phe Lys Arg Arg Asn Gln Glu Arg Leu Asn Pro Arg Asp 150 155 160	956
GTG GAG TAC GGT ACT ATC GAA GGG CTC ATC ACC ACC AAC GTC GGA GAT Val Glu Tyr Gly Thr Ile Glu Gly Leu Ile Thr Thr Asn Val Gly Asp 165 170 175	1004
AGC ACT CTA GCG GAA TTA CTA GAT CAC TCA TGT ACA TCA GGA AGT GGC Ser Thr Leu Ala Glu Leu Leu Asp His Ser Cys Thr Ser Gly Ser Gly 180 185 190	1052
TCC GGT CTT CCT TTT CTG GTA CAG AGA ACT GTG GCT CGA CAG ATA ACC Ser Gly Leu Pro Phe Leu Val Gln Arg Thr Val Ala Arg Gln Ile Thr 195 200 205	1100
CTG TTG GAG TGT GTC GGG AAG GGC CGG TAT GGA GAA GTG TGG AGG GGC Leu Leu Glu Cys Val Gly Lys Gly Arg Tyr Gly Glu Val Trp Arg Gly 210 215 220 225	1148
AGC TGG CAA GGC GAA AAT GTT GCT GTG AAG ATC TTC TCC TCC CGT GAT Ser Trp Gln Gly Glu Asn Val Ala Val Lys Ile Phe Ser Ser Arg Asp 230 235 240	1196
GAG AAG TCG TGG TTC AGG GAG ACA GAA TTG TAC AAC ACG GTG ATG CTG Glu Lys Ser Trp Phe Arg Glu Thr Glu Leu Tyr Asn Thr Val Met Leu 245 250 255	1244
AGG CAT GAG AAT ATC TTA GGT TTC ATT GCT TCA GAC ATG ACC TCT AGA Arg His Glu Asn Ile Leu Gly Phe Ile Ala Ser Asp Met Thr Ser Arg 260 265 270	1292
CAC TCC AGT ACC CAG CTG TGG CTC ATT ACA CAT TAC CAC GAA ATG GGA His Ser Ser Thr Gln Leu Trp Leu Ile Thr His Tyr His Glu Met Gly 275 280 285	1340
TCG TTG TAT GAC TAC CTT CAG CTC ACC ACT CTG GAC ACC GTT AGC TGC Ser Leu Tyr Asp Tyr Leu Gln Leu Thr Thr Leu Asp Thr Val Ser Cys 290 295 300 305	1388
CTT CGG ATC GTG TTG TCC ATA GCC AGC GGC CTT GCA CAC TTG CAC ATA Leu Arg Ile Val Leu Ser Ile Ala Ser Gly Leu Ala His Leu His Ile 310 315 320	1436
GAG ATA TTT GGG ACC CAG GGG AAG TCT GCC ATC GCC CAC CGA GAT CTA Glu Ile Phe Gly Thr Gln Gly Lys Ser Ala Ile Ala His Arg Asp Leu 325 330 335	1484
AAG AGC AAA AAC ATC CTC GTG AAG AAG AAC GGA CAG TGC TGC ATA GCA Lys Ser Lys Asn Ile Leu Val Lys Lys Asn Gly Gln Cys Cys Ile Ala 340 345 350	1532
GAT TTG GGC CTG GCA GTC ATG CAT TCC CAG AGC ACG AAT CAG CTT GAT Asp Leu Gly Leu Ala Val Met His Ser Gln Ser Thr Asn Gln Leu Asp 355 360 365	1580
GTG GGA AAC AAC CCC CGT GTG GGG ACC AAG CGC TAC ATG GCC CCT GAA Val Gly Asn Asn Pro Arg Val Gly Thr Lys Arg Tyr Met Ala Pro Glu 370 375 380 385	1628
GTG CTT GAT GAA ACC ATC CAA GTG GAT TGC TTT GAT TCT TAT AAG AGG Val Leu Asp Glu Thr Ile Gln Val Asp Cys Phe Asp Ser Tyr Lys Arg 390 395 400	1676
GTC GAT ATT TGG GCC TTT GGC CTC GTT CTG TGG GAA GTG GCC AGG AGG Val Asp Ile Trp Ala Phe Gly Leu Val Leu Trp Glu Val Ala Arg Arg 405 410 415	1724

ATG GTG AGC AAT GGT ATA GTG GAA GAT TAC AAG CCA CCA TTC TAT GAT Met Val Ser Asn Gly Ile Val Glu Asp Tyr Lys Pro Pro Phe Tyr Asp 420 425 430	1772
GTT GTT CCC AAT GAC CCA AGT TTT GAA GAT ATG AGG AAA GTT GTC TGT Val Val Pro Asn Asp Pro Ser Phe Glu Asp Met Arg Lys Val Val Cys 435 440 445	1820
GTG GAT CAA CAG AGG CCA AAC ATA CCT AAC AGA TGG TTC TCA GAC CCG Val Asp Gln Gln Arg Pro Asn Ile Pro Asn Arg Trp Phe Ser Asp Pro 450 455 460 465	1868
ACA TTA ACT TCT CTG GCG AAG CTG ATG AAA GAA TGC TGG TAC CAG AAC Thr Leu Thr Ser Leu Ala Lys Leu Met Lys Glu Cys Trp Tyr Gln Asn 470 475 480	1916
CCA TCC GCC AGA CTC ACA GCT CTA CGT ATC AAA AAG ACT TTG ACC AAA Pro Ser Ala Arg Leu Thr Ala Leu Arg Ile Lys Lys Thr Leu Thr Lys 485 490 495	1964
ATT GAT AAC TCC CTA GAC AAA TTA AAA ACT GAC TGT TGACATTGTC Ile Asp Asn Ser Leu Asp Lys Leu Lys Thr Asp Cys 500 505	2010
ACCGGTGTCA AGAAGGAGAG TCAATGCTGT CATTGTCCAG CTGGGACCTA ATGCTGGCCT	2070
GACTGGTTGT CAGAACAGAA TCCATCTGTC CCCCTCTCCC CCCAACTCCC GAAGTGGCTG	2130
CTTTGACAAA AGCAGATGTC TCTTCCCAGC CATGTTCCGG GGGAGACACC AAAACCACCC	2190
TAACCTCGCT CAGAACTGT GACTCGAGCA CTTGATGAAC TGTTACACACC GCAAAGACTA	2250
ACGGTGGGCA GGTATGTTTG CAAGGGGGAG GGAAGTGGAG GAGCACAGAG AGATCCTGCA	2310
GGAGATCTGG GCATTAGGAC AGTGGCTCTT TCGGTATCTT CCACGGGTCT CCTAGACTCG	2370
<del>CCCCACGGGA</del> AACTCAAGGA GGCGGTGAAT TCGTAATCAG CAATATTGGC TGCGCCTACT	2430
CTTCTCTGTT GCACTAGGAA TTCTCTGCAT TCCTTACTTG CACTGTCGTC CTTAATCTTA	2490
AAGACCCGAC TTGCCAAAAC ATTGGCTGCC TACTTCACTG GCCTGTCTCT GGACAATAGG	2550
AATTCAATCT GCGGAAACAA AAATGTAATG TTGGACTTTG CTGCATTTTA CACACGTGCC	2610
GATGTTTACA ACGATGCAAA CATTAGGAAT TGTTTAGACA CAACTTTGCA AATTATTTAT	2670
TACTGGTGCA CTTAGCAGTT TTTGTTTTTT TTTGTTTTTT TGTTTTTTTT TTGTTTTGTT	2730
TTGTTTTTAT ATATAAACT GCCTCGTGCG TATGTTAAAG CTTATTTTTTA TGTGGTCTTA	2790
TGATTTTATT ACCGAAATGT TTTTAACACC CGATTCTGAA ATGGATGTTT TCTTTTATTA	2850
TCAGTTAAAT TCACATTTTA AATGCTTCAC TTTTTTTTTTA TGTGTGTAGA CTGTAACTTT	2910
CTTTTCAGTT AGTATACAGA ACGTATTTAG CCATTACCCA TGCAACACCA CCCAATATAT	2970
TACTGATTTA GAAGCAAAGA TTTCAGTAGA ATTTTAGTCC CAAACGCTGT GGGGGGGAAA	3030
TGCATCTTCT TCGGAACTAT CCATTACATG CATTTAACT CTGCCAGAAA AAAAAATAAC	3090
TATTTTGTTT TAATCTACTT TTTGTATTTA GTAGTTATTT GTATAAATTA AATAAACTGT	3150
TTTCAAGTCA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAT	3210
AAAAAAAAAA AAAGCGGCCG CAGAATTC	3238

## (2) INFORMATION FOR SEQ ID NO:6:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 509 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

```

Met Val Asp Gly Ala Met Ile Leu Ser Val Leu Met Met Met Ala Leu
 1      5      10      15
Pro Ser Pro Ser Met Glu Asp Glu Glu Pro Lys Val Asn Pro Lys Leu
 20      25      30
Tyr Met Cys Val Cys Glu Gly Leu Ser Cys Gly Asn Glu Asp His Cys
 35      40      45
Glu Gly Gln Gln Cys Phe Ser Ser Leu Ser Val Asn Asp Gly Phe Arg
 50      55      60
Val Tyr Gln Lys Gly Cys Phe Gln Val Tyr Glu Gln Gly Lys Met Thr
 65      70      75      80
Cys Lys Thr Pro Pro Ser Pro Gly Gln Ala Val Glu Cys Cys Gln Gly
 85      90      95
Asp Trp Cys Asn Arg Asn Val Thr Ala Arg Leu Pro Thr Lys Gly Lys
100      105      110
Ser Phe Pro Gly Ser Gln Asn Phe His Leu Glu Val Gly Leu Ile Ile
115      120      125
Leu Ser Val Val Phe Ala Val Cys Leu Phe Ala Cys Ile Leu Gly Val
130      135      140
Ala Leu Arg Lys Phe Lys Arg Arg Asn Gln Glu Arg Leu Asn Pro Arg
145      150      155      160
Asp Val Glu Tyr Gly Thr Ile Glu Gly Leu Ile Thr Thr Asn Val Gly
165      170      175
Asp Ser Thr Leu Ala Glu Leu Leu Asp His Ser Cys Thr Ser Gly Ser
180      185      190
Gly Ser Gly Leu Pro Phe Leu Val Gln Arg Thr Val Ala Arg Gln Ile
195      200      205
Thr Leu Leu Glu Cys Val Gly Lys Gly Arg Tyr Gly Glu Val Trp Arg
210      215      220
Gly Ser Trp Gln Gly Glu Asn Val Ala Val Lys Ile Phe Ser Ser Arg
225      230      235      240
Asp Glu Lys Ser Trp Phe Arg Glu Thr Glu Leu Tyr Asn Thr Val Met
245      250      255
Leu Arg His Glu Asn Ile Leu Gly Phe Ile Ala Ser Asp Met Thr Ser
260      265      270
Arg His Ser Ser Thr Gln Leu Trp Leu Ile Thr His Tyr His Glu Met
275      280      285
Gly Ser Leu Tyr Asp Tyr Leu Gln Leu Thr Thr Leu Asp Thr Val Ser
290      295      300

```

Cys Leu Arg Ile Val Leu Ser Ile Ala Ser Gly Leu Ala His Leu His  
 305 310 315 320  
 Ile Glu Ile Phe Gly Thr Gln Gly Lys Ser Ala Ile Ala His Arg Asp  
 325 330 335  
 Leu Lys Ser Lys Asn Ile Leu Val Lys Lys Asn Gly Gln Cys Cys Ile  
 340 345 350  
 Ala Asp Leu Gly Leu Ala Val Met His Ser Gln Ser Thr Asn Gln Leu  
 355 360 365  
~~Asp Val Gly Asn Asn Pro Arg Val Gly Thr Lys Arg Tyr Met Ala Pro~~  
~~370 375 380~~  
 Glu Val Leu Asp Glu Thr Ile Gln Val Asp Cys Phe Asp Ser Tyr Lys  
 385 390 395 400  
 Arg Val Asp Ile Trp Ala Phe Gly Leu Val Leu Trp Glu Val Ala Arg  
 405 410 415  
 Arg Met Val Ser Asn Gly Ile Val Glu Asp Tyr Lys Pro Pro Phe Tyr  
 420 425 430  
 Asp Val Val Pro Asn Asp Pro Ser Phe Glu Asp Met Arg Lys Val Val  
 435 440 445  
 Cys Val Asp Gln Gln Arg Pro Asn Ile Pro Asn Arg Trp Phe Ser Asp  
 450 455 460  
 Pro Thr Leu Thr Ser Leu Ala Lys Leu Met Lys Glu Cys Trp Tyr Gln  
 465 470 475 480  
 Asn Pro Ser Ala Arg Leu Thr Ala Leu Arg Ile Lys Lys Thr Leu Thr  
 485 490 495  
~~Lys Ile Asp Asn Ser Leu Asp Lys Leu Lys Thr Asp Cys~~  
~~500 505~~

## (2) INFORMATION FOR SEQ ID NO:7:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1647 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (vii) IMMEDIATE SOURCE:

- (B) CLONE: W-101

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 80..1594

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GAATCTGCGG CCGCGAGGGA GAGAGGCGCC GGGGGCGCGC GCGCGCGCTG GGCGCTGCTG 60  
 GGCTGCGGCG GCGGTTACT ATG GCG GAG TCG GCC GGA GCC TCC TCC TTC TTC 112  
 Met Ala Glu Ser Ala Gly Ala Ser Ser Phe Phe  
 1 5 10

CCC	CTT	GTT	GTC	CTC	CTG	CTC	GCC	GGC	AGC	GGC	GGG	TCC	GGG	CCC	CGG	160
Pro	Leu	Val	Val	Leu	Leu	Leu	Ala	Gly	Ser	Gly	Gly	Ser	Gly	Pro	Arg	
			15					20					25			
GGG	ATC	CAG	GCT	CTG	CTG	TGT	GCG	TGC	ACC	AGC	TGC	CTA	CAG	ACC	AAC	208
Gly	Ile	Gln	Ala	Leu	Leu	Cys	Ala	Cys	Thr	Ser	Cys	Leu	Gln	Thr	Asn	
		30					35					40				
TAC	ACC	TGT	GAG	ACA	GAT	GGG	GCT	TGC	ATG	GTC	TCC	ATC	TTT	AAC	CTG	256
Tyr	Thr	Cys	Glu	Thr	Asp	Gly	Ala	Cys	Met	Val	Ser	Ile	Phe	Asn	Leu	
	45					50					55					
GAT	GGC	GTG	GAG	CAC	CAT	GTA	CGT	ACC	TGC	ATC	CCC	AAG	GTG	GAG	CTG	304
Asp	Gly	Val	Glu	His	His	Val	Arg	Thr	Cys	Ile	Pro	Lys	Val	Glu	Leu	
60					65					70					75	
GTT	CCT	GCT	GGA	AAG	CCC	TTC	TAC	TGC	CTG	AGT	TCA	GAG	GAT	CTG	CGC	352
Val	Pro	Ala	Gly	Lys	Pro	Phe	Tyr	Cys	Leu	Ser	Ser	Glu	Asp	Leu	Arg	
				80					85					90		
AAC	ACA	CAC	TGC	TGC	TAT	ATT	GAC	TTC	TGC	AAC	AAG	ATT	GAC	CTC	AGG	400
Asn	Thr	His	Cys	Cys	Tyr	Ile	Asp	Phe	Cys	Asn	Lys	Ile	Asp	Leu	Arg	
			95					100					105			
GTC	CCC	AGC	GGA	CAC	CTC	AAG	GAG	CCT	GCG	CAC	CCC	TCC	ATG	TGG	GGC	448
Val	Pro	Ser	Gly	His	Leu	Lys	Glu	Pro	Ala	His	Pro	Ser	Met	Trp	Gly	
		110					115					120				
CCT	GTG	GAG	CTG	GTC	GGC	ATC	ATC	GCC	GGC	CCC	GTC	TTC	CTC	CTC	TTC	496
Pro	Val	Glu	Leu	Val	Gly	Ile	Ile	Ala	Gly	Pro	Val	Phe	Leu	Leu	Phe	
	125					130					135					
CTT	ATC	ATT	ATC	ATC	GTC	TTC	CTG	GTC	ATC	AAC	TAT	CAC	CAG	CGT	GTC	544
Leu	Ile	Ile	Ile	Ile	Val	Phe	Leu	Val	Ile	Asn	Tyr	His	Gln	Arg	Val	
140					145					150					155	
TAC	CAT	AAC	CGC	CAG	AGG	TTG	GAC	ATG	GAG	GAC	CCC	TCT	TGC	GAG	ATG	592
Tyr	His	Asn	Arg	Gln	Arg	Leu	Asp	Met	Glu	Asp	Pro	Ser	Cys	Glu	Met	
				160					165					170		
TGT	CTC	TCC	AAA	GAC	AAG	ACG	CTC	CAG	GAT	CTC	GTC	TAC	GAC	CTC	TCC	640
Cys	Leu	Ser	Lys	Asp	Lys	Thr	Leu	Gln	Asp	Leu	Val	Tyr	Asp	Leu	Ser	
			175					180					185			
ACG	TCA	GGG	TCT	GGC	TCA	GGG	TTA	CCC	CTT	TTT	GTC	CAG	CGC	ACA	GTG	688
Thr	Ser	Gly	Ser	Gly	Ser	Gly	Leu	Pro	Leu	Phe	Val	Gln	Arg	Thr	Val	
		190					195					200				
GCC	CGA	ACC	ATT	GTT	TTA	CAA	GAG	ATT	ATC	GGC	AAG	GGC	CGG	TTC	GGG	736
Ala	Arg	Thr	Ile	Val	Leu	Gln	Glu	Ile	Ile	Gly	Lys	Gly	Arg	Phe	Gly	
	205					210					215					
GAA	GTA	TGG	CGT	GGT	CGC	TGG	AGG	GGT	GGT	GAC	GTG	GCT	GTG	AAA	ATC	784
Glu	Val	Trp	Arg	Gly	Arg	Trp	Arg	Gly	Gly	Asp	Val	Ala	Val	Lys	Ile	
220					225					230					235	
TTC	TCT	TCT	CGT	GAA	GAA	CGG	TCT	TGG	TTC	CGT	GAA	GCA	GAG	ATC	TAC	832
Phe	Ser	Ser	Arg	Glu	Glu	Arg	Ser	Trp	Phe	Arg	Glu	Ala	Glu	Ile	Tyr	
				240					245					250		
CAG	ACC	GTC	ATG	CTG	CGC	CAT	GAA	AAC	ATC	CTT	GGC	TTT	ATT	GCT	GCT	880
Gln	Thr	Val	Met	Leu	Arg	His	Glu	Asn	Ile	Leu	Gly	Phe	Ile	Ala	Ala	
			255					260					265			
GAC	AAT	AAA	GAT	AAT	GGC	ACC	TGG	ACC	CAG	CTG	TGG	CTT	GTC	TCT	GAC	928
Asp	Asn	Lys	Asp	Asn	Gly	Thr	Trp	Thr	Gln	Leu	Trp	Leu	Val	Ser	Asp	
		270					275					280				



TAT CAC GAG CAT GGC TCA CTG TTT GAT TAT CTG AAC CGC TAC ACC GTG Tyr His Glu His Gly Ser Leu Phe Asp Tyr Leu Asn Arg Tyr Thr Val 285 290 295	976
ACC ATT GAG GGC ATG ATT AAG CTA GCC TTG TCT GCA GCC AGT GGT TTG Thr Ile Glu Gly Met Ile Lys Leu Ala Leu Ser Ala Ala Ser Gly Leu 300 305 310 315	1024
GCA CAC CTG CAT ATG GAG ATT GTG GGC ACT CAA GGG AAG CCG GGA ATT Ala His Leu His Met Glu Ile Val Gly Thr Gln Gly Lys Pro Gly Ile 320 325 330	1072
GCT CAT CGA GAC TTG AAG TCA AAG AAC ATC CTG GTG AAA AAA AAT GGC Ala His Arg Asp Leu Lys Ser Lys Asn Ile Leu Val Lys Lys Asn Gly 335 340 345	1120
ATG TGT GCC ATT GCA GAC CTG GGC CTG GCT GTC CGT CAT GAT GCG GTC Met Cys Ala Ile Ala Asp Leu Gly Leu Ala Val Arg His Asp Ala Val 350 355 360	1168
ACT GAC ACC ATA GAC ATT GCT CCA AAT CAG AGG GTG GGG ACC AAA CGA Thr Asp Thr Ile Asp Ile Ala Pro Asn Gln Arg Val Gly Thr Lys Arg 365 370 375	1216
TAC ATG GCT CCT GAA GTC CTT GAC GAG ACA ATC AAC ATG AAG CAC TTT Tyr Met Ala Pro Glu Val Leu Asp Glu Thr Ile Asn Met Lys His Phe 380 385 390 395	1264
GAC TCC TTC AAA TGT GCC GAC ATC TAT GCC CTC GGG CTT GTC TAC TGG Asp Ser Phe Lys Cys Ala Asp Ile Tyr Ala Leu Gly Leu Val Tyr Trp 400 405 410	1312
GAG ATT GCA CGA AGA TGC AAT TCT GGA GGA GTC CAT GAA GAC TAT CAA Glu Ile Ala Arg Arg Cys Asn Ser Gly Gly Val His Glu Asp Tyr Gln 415 420 425	1360
CTG CCG TAT TAC GAC TTA GTG CCC TCC GAC CCT TCC ATT GAG GAG ATG Leu Pro Tyr Tyr Asp Leu Val Pro Ser Asp Pro Ser Ile Glu Glu Met 430 435 440	1408
CGA AAG GTT GTA TGT GAC CAG AAG CTA CGG CCC AAT GTC CCC AAC TGG Arg Lys Val Val Cys Asp Gln Lys Leu Arg Pro Asn Val Pro Asn Trp 445 450 455	1456
TGG CAG AGT TAT GAG GCC TTG CGA GTG ATG GGA AAG ATG ATG CGG GAG Trp Gln Ser Tyr Glu Ala Leu Arg Val Met Gly Lys Met Met Arg Glu 460 465 470 475	1504
TGC TGG TAC GCC AAT GGT GCT GCC CGT CTG ACA GCT CTG CGC ATC AAG Cys Trp Tyr Ala Asn Gly Ala Ala Arg Leu Thr Ala Leu Arg Ile Lys 480 485 490	1552
AAG ACT CTG TCC CAG CTA AGC GTG CAG GAA GAT GTG AAG ATT Lys Thr Leu Ser Gln Leu Ser Val Gln Glu Asp Val Lys Ile 495 500 505	1594
TAAGCTGTTA AGATGCCTAC ACAAAGAACC TGGGCAGTGA GGATGACTGC AGG	1647

## (2) INFORMATION FOR SEQ ID NO:8:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 505 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Met Ala Glu Ser Ala Gly Ala Ser Ser Phe Phe Pro Leu Val Val Leu  
 1 5 10 15  
 Leu Leu Ala Gly Ser Gly Gly Ser Gly Pro Arg Gly Ile Gln Ala Leu  
 20 25 30  
 Leu Cys Ala Cys Thr Ser Cys Leu Gln Thr Asn Tyr Thr Cys Glu Thr  
 35 40 45  
 Asp Gly Ala Cys Met Val Ser Ile Phe Asn Leu Asp Gly Val Glu His  
 50 55 60  
 His Val Arg Thr Cys Ile Pro Lys Val Glu Leu Val Pro Ala Gly Lys  
 65 70 75 80  
 Pro Phe Tyr Cys Leu Ser Ser Glu Asp Leu Arg Asn Thr His Cys Cys  
 85 90 95  
 Tyr Ile Asp Phe Cys Asn Lys Ile Asp Leu Arg Val Pro Ser Gly His  
 100 105 110  
 Leu Lys Glu Pro Ala His Pro Ser Met Trp Gly Pro Val Glu Leu Val  
 115 120 125  
 Gly Ile Ile Ala Gly Pro Val Phe Leu Leu Phe Leu Ile Ile Ile Ile  
 130 135 140  
 Val Phe Leu Val Ile Asn Tyr His Gln Arg Val Tyr His Asn Arg Gln  
 145 150 155 160  
 Arg Leu Asp Met Glu Asp Pro Ser Cys Glu Met Cys Leu Ser Lys Asp  
 165 170 175  
 Lys Thr Leu Gln Asp Leu Val Tyr Asp Leu Ser Thr Ser Gly Ser Gly  
 180 185 190  
 Ser Gly Leu Pro Leu Phe Val Gln Arg Thr Val Ala Arg Thr Ile Val  
 195 200 205  
 Leu Gln Glu Ile Ile Gly Lys Gly Arg Phe Gly Glu Val Trp Arg Gly  
 210 215 220  
 Arg Trp Arg Gly Gly Asp Val Ala Val Lys Ile Phe Ser Ser Arg Glu  
 225 230 235 240  
 Glu Arg Ser Trp Phe Arg Glu Ala Glu Ile Tyr Gln Thr Val Met Leu  
 245 250 255  
 Arg His Glu Asn Ile Leu Gly Phe Ile Ala Ala Asp Asn Lys Asp Asn  
 260 265 270  
 Gly Thr Trp Thr Gln Leu Trp Leu Val Ser Asp Tyr His Glu His Gly  
 275 280 285  
 Ser Leu Phe Asp Tyr Leu Asn Arg Tyr Thr Val Thr Ile Glu Gly Met  
 290 295 300  
 Ile Lys Leu Ala Leu Ser Ala Ala Ser Gly Leu Ala His Leu His Met  
 305 310 315 320  
 Glu Ile Val Gly Thr Gln Gly Lys Pro Gly Ile Ala His Arg Asp Leu  
 325 330 335  
 Lys Ser Lys Asn Ile Leu Val Lys Lys Asn Gly Met Cys Ala Ile Ala  
 340 345 350

Asp Leu Gly Leu Ala Val Arg His Asp Ala Val Thr Asp Thr Ile Asp  
 355 360 365  
 Ile Ala Pro Asn Gln Arg Val Gly Thr Lys Arg Tyr Met Ala Pro Glu  
 370 375 380  
 Val Leu Asp Glu Thr Ile Asn Met Lys His Phe Asp Ser Phe Lys Cys  
 385 390 395 400  
 Ala Asp Ile Tyr Ala Leu Gly Leu Val Tyr Trp Glu Ile Ala Arg Arg  
 405 410 415  
 Cys Asn-Ser Gly Gly Val His-Glu Asp Tyr Gln-Leu Pro Tyr Tyr Asp  
 420 425 430  
 Leu Val Pro Ser Asp Pro Ser Ile Glu Glu Met Arg Lys Val Val Cys  
 435 440 445  
 Asp Gln Lys Leu Arg Pro Asn Val Pro Asn Trp Trp Gln Ser Tyr Glu  
 450 455 460  
 Ala Leu Arg Val Met Gly Lys Met Met Arg Glu Cys Trp Tyr Ala Asn  
 465 470 475 480  
 Gly Ala Ala Arg Leu Thr Ala Leu Arg Ile Lys Lys Thr Leu Ser Gln  
 485 490 495  
 Leu Ser Val Gln Glu Asp Val Lys Ile  
 500 505

## (2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1794 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vii) IMMEDIATE SOURCE:

- (B) CLONE: W-120

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 83..1591

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GAATTCGCGG CCGCGGGCGA GGCTTCCTGA GGAGAAGCTG CGGCCGGGGC CGGGCCGGGC	60
CACAAACAGT GCGGCGGGA-CC-ATG-GAG GCG GCG-GCC GCT GCT CCA CGT CGT	112
Met Glu Ala Ala Ala Ala Ala Pro Arg Arg	
1 5 10	
CCG CAG CTC CTC ATC GTG TTG GTG GCG GCG GCG ACG CTG CTC CCG GGG	160
Pro Gln Leu Leu Ile Val Leu Val Ala Ala Ala Thr Leu Leu Pro Gly	
15 20 25	
GCG AAG GCA TTA CAG TGT TTC TGC CAC CTC TGT ACA AAG GAT AAT TTT	208
Ala Lys Ala Leu Gln Cys Phe Cys His Leu Cys Thr Lys Asp Asn Phe	
30 35 40	

ACC	TGT	GAG	ACA	GAT	GGT	CTT	TGC	TTT	GTC	TCA	GTC	ACT	GAG	ACC	ACA	256
Thr	Cys	Glu	Thr	Asp	Gly	Leu	Cys	Phe	Val	Ser	Val	Thr	Glu	Thr	Thr	
		45					50					55				
GAC	AAA	GTT	ATA	CAC	AAT	AGT	ATG	TGT	ATA	GCT	GAA	ATT	GAC	CTA	ATT	304
Asp	Lys	Val	Ile	His	Asn	Ser	Met	Cys	Ile	Ala	Glu	Ile	Asp	Leu	Ile	
	60					65					70					
CCT	CGA	GAC	AGG	CCA	TTT	GTA	TGT	GCA	CCA	TCT	TCA	AAA	ACA	GGG	GCA	352
Pro	Arg	Asp	Arg	Pro	Phe	Val	Cys	Ala	Pro	Ser	Ser	Lys	Thr	Gly	Ala	
	75				80					85					90	
GTT	ACT	ACA	ACA	TAT	TGC	TGC	AAT	CAG	GAC	CAC	TGC	AAT	AAA	ATA	GAA	400
Val	Thr	Thr	Thr	Tyr	Cys	Cys	Asn	Gln	Asp	His	Cys	Asn	Lys	Ile	Glu	
				95					100					105		
CTC	CCA	ACT	ACA	GGA	CCT	TTT	TCA	GAA	AAG	CAG	TCA	GCT	GGC	CTT	GGT	448
Leu	Pro	Thr	Thr	Gly	Pro	Phe	Ser	Glu	Lys	Gln	Ser	Ala	Gly	Leu	Gly	
			110					115					120			
CCT	GTG	GAG	CTG	GCA	GCT	GTC	ATT	GCT	GGT	CCA	GTC	TGC	TTC	GTC	TGC	496
Pro	Val	Glu	Leu	Ala	Ala	Val	Ile	Ala	Gly	Pro	Val	Cys	Phe	Val	Cys	
		125					130					135				
ATT	GCA	CTT	ATG	CTG	ATG	GTC	TAT	ATC	TGC	CAT	AAC	CGC	ACT	GTC	ATT	544
Ile	Ala	Leu	Met	Leu	Met	Val	Tyr	Ile	Cys	His	Asn	Arg	Thr	Val	Ile	
	140					145					150					
CAC	CAC	CGT	GTG	CCA	AAT	GAA	GAG	GAT	CCA	TCA	CTA	GAT	CGC	CCT	TTC	592
His	His	Arg	Val	Pro	Asn	Glu	Glu	Asp	Pro	Ser	Leu	Asp	Arg	Pro	Phe	
	155				160					165					170	
ATT	TCA	GAG	GGC	ACC	ACC	TTA	AAA	GAT	TTA	ATT	TAT	GAT	ATG	ACA	ACA	640
Ile	Ser	Glu	Gly	Thr	Thr	Leu	Lys	Asp	Leu	Ile	Tyr	Asp	Met	Thr	Thr	
				175					180					185		
TCA	GGG	TCT	GGA	TCA	GGT	TTA	CCA	CTG	CTT	GTT	CAA	AGA	ACA	ATT	GCC	688
Ser	Gly	Ser	Gly	Ser	Gly	Leu	Pro	Leu	Leu	Val	Gln	Arg	Thr	Ile	Ala	
			190					195					200			
AGG	ACC	ATT	GTG	TTA	CAA	GAA	AGC	ATT	GGC	AAA	GGT	CGG	TTT	GGA	GAA	736
Arg	Thr	Ile	Val	Leu	Gln	Glu	Ser	Ile	Gly	Lys	Gly	Arg	Phe	Gly	Glu	
		205					210					215				
GTT	TGG	CGA	GGC	AAA	TGG	CGG	GGA	GAA	GAA	GTT	GCT	GTG	AAG	ATA	TTC	784
Val	Trp	Arg	Gly	Lys	Trp	Arg	Gly	Glu	Glu	Val	Ala	Val	Lys	Ile	Phe	
	220					225					230					
TCT	TCT	AGA	GAA	GAG	CGT	TCA	TGG	TTC	CGA	GAG	GCA	GAG	ATT	TAT	CAG	832
Ser	Ser	Arg	Glu	Glu	Arg	Ser	Trp	Phe	Arg	Glu	Ala	Glu	Ile	Tyr	Gln	
	235				240					245					250	
ACT	GTA	ATG	TTA	CGC	CAT	GAA	AAT	ATC	CTG	GGA	TTT	ATA	GCA	GCA	GAC	880
Thr	Val	Met	Leu	Arg	His	Glu	Asn	Ile	Leu	Gly	Phe	Ile	Ala	Ala	Asp	
				255					260					265		
AAC	AAA	GAC	AAT	GGG	ACA	TGG	ACG	CAG	CTG	TGG	TTG	GTG	TCA	GAT	TAT	928
Asn	Lys	Asp	Asn	Gly	Thr	Trp	Thr	Gln	Leu	Trp	Leu	Val	Ser	Asp	Tyr	
			270					275					280			
CAT	GAG	CAT	GGA	TCC	CTT	TTC	GAT	TAC	TTG	AAT	AGA	TAC	ACT	GTT	ACT	976
His	Glu	His	Gly	Ser	Leu	Phe	Asp	Tyr	Leu	Asn	Arg	Tyr	Thr	Val	Thr	
		285					290					295				
GTG	GAA	GGA	ATG	ATC	AAG	CTT	GCT	CTG	TCC	ACA	GCA	AGT	GGT	CTT	GCC	1024
Val	Glu	Gly	Met	Ile	Lys	Leu	Ala	Leu	Ser	Thr	Ala	Ser	Gly	Leu	Ala	
	300					305					310					

CAT CTT CAC ATG GAG ATT GTT GGT ACC CAA GGA AAA CCA GCT ATT GCC His Leu His Met Glu Ile Val Gly Thr Gln Gly Lys Pro Ala Ile Ala 315 320 325 330	1072
CAT AGA GAT TTG AAA TCA AAG AAT ATC TTG GTG AAG AAA AAT GGA ACC His Arg Asp Leu Lys Ser Lys Asn Ile Leu Val Lys Lys Asn Gly Thr 335 340 345	1120
TGT TGT ATT GCA GAC TTG GGA CTT GCT GTG AGA CAT GAT TCT GCC ACA Cys Cys Ile Ala Asp Leu Gly Leu Ala Val Arg His Asp Ser Ala Thr 350 355 360	1168
GAT ACA ATT GAT ATT GCT CCA AAC CAC AGA GTA GGC ACT AAA AGG TAC Asp Thr Ile Asp Ile Ala Pro Asn His Arg Val Gly Thr Lys Arg Tyr 365 370 375	1216
ATG GCC CCT GAA GTT CTA GAT GAT TCC ATA AAT ATG AAA CAT TTT GAA Met Ala Pro Glu Val Leu Asp Asp Ser Ile Asn Met Lys His Phe Glu 380 385 390	1264
TCC TTC AAA CGC GCT GAC ATC TAT GCA ATG GGC TTA GTG TTC TGG GAA Ser Phe Lys Arg Ala Asp Ile Tyr Ala Met Gly Leu Val Phe Trp Glu 395 400 405 410	1312
ATT GCT CGA CGC TGT TCT ATT GGT GGA ATC CAT GAA GAC TAT CAG TTG Ile Ala Arg Arg Cys Ser Ile Gly Gly Ile His Glu Asp Tyr Gln Leu 415 420 425	1360
CCT TAT TAT GAT CTT GTA CCT TCT GAT CCA TCG GTT GAA GAA ATG AGA Pro Tyr Tyr Asp Leu Val Pro Ser Asp Pro Ser Val Glu Glu Met Arg 430 435 440	1408
AAA GTA GTT TGC GAA CAG AAG TTA AGG CCA AAT ATT CCA AAC AGA TGG Lys Val Val Cys Glu Gln Lys Leu Arg Pro Asn Ile Pro Asn Arg Trp 445 450 455	1456
CAG AGC TGT GAG GCC TTG AGA GTG ATG GCT AAA ATT ATG AGA GAA TGC Gln Ser Cys Glu Ala Leu Arg Val Met Ala Lys Ile Met Arg Glu Cys 460 465 470	1504
TGG TAT GCC AAT GGA GCA GCA AGG CTG ACA GCT TTG CGA ATT AAA AAA Trp Tyr Ala Asn Gly Ala Ala Arg Leu Thr Ala Leu Arg Ile Lys Lys 475 480 485 490	1552
ACA TTG TCA CAA CTC AGC CAA CAG GAA GGC ATC AAA ATG TAACTGAAAC Thr Leu Ser Gln Leu Ser Gln Gln Glu Gly Ile Lys Met 495 500	1601
ACCGTGGGAA CTCTGCTCTC TTCATATCTG CTCCTGGGTG TTTAGGAGGC TGGTTGTTCT	1661
ACCTCACTGA GAGAACAGAG GGCTCTGCTT CCTCTTGCAG CAGTGAATA TGGTCAACTG	1721
AAAGCTTCCC AGGGTTTCTC TGGGCCCAGA GGCAGCCGTG GGGTCCTTCT GTGCACTATG	1781
GATAACTTCT TCC	1794

## (2) INFORMATION FOR SEQ ID NO:10:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 503 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:



Met Glu Ala Ala Ala Ala Ala Pro Arg Arg Pro Gln Leu Leu Ile Val  
 1 5 10 15  
 Leu Val Ala Ala Ala Thr Leu Leu Pro Gly Ala Lys Ala Leu Gln Cys  
 20 25 30  
 Phe Cys His Leu Cys Thr Lys Asp Asn Phe Thr Cys Glu Thr Asp Gly  
 35 40 45  
 Leu Cys Phe Val Ser Val Thr Glu Thr Thr Asp Lys Val Ile His Asn  
 50 55 60  
~~Ser Met Cys Ile Ala Glu Ile Asp Leu Ile Pro Arg Asp Arg Pro Phe~~  
 65 70 75 80  
 Val Cys Ala Pro Ser Ser Lys Thr Gly Ala Val Thr Thr Thr Tyr Cys  
 85 90 95  
 Cys Asn Gln Asp His Cys Asn Lys Ile Glu Leu Pro Thr Thr Gly Pro  
 100 105 110  
 Phe Ser Glu Lys Gln Ser Ala Gly Leu Gly Pro Val Glu Leu Ala Ala  
 115 120 125  
 Val Ile Ala Gly Pro Val Cys Phe Val Cys Ile Ala Leu Met Leu Met  
 130 135 140  
 Val Tyr Ile Cys His Asn Arg Thr Val Ile His His Arg Val Pro Asn  
 145 150 155 160  
 Glu Glu Asp Pro Ser Leu Asp Arg Pro Phe Ile Ser Glu Gly Thr Thr  
 165 170 175  
 Leu Lys Asp Leu Ile Tyr Asp Met Thr Thr Ser Gly Ser Gly Ser Gly  
 180 185 190  
 Leu Pro Leu Leu Val Gln Arg Thr Ile Ala Arg Thr Ile Val Leu Gln  
 195 200 205  
 Glu Ser Ile Gly Lys Gly Arg Phe Gly Glu Val Trp Arg Gly Lys Trp  
 210 215 220  
 Arg Gly Glu Glu Val Ala Val Lys Ile Phe Ser Ser Arg Glu Glu Arg  
 225 230 235 240  
 Ser Trp Phe Arg Glu Ala Glu Ile Tyr Gln Thr Val Met Leu Arg His  
 245 250 255  
 Glu Asn Ile Leu Gly Phe Ile Ala Ala Asp Asn Lys Asp Asn Gly Thr  
 260 265 270  
 Trp Thr Gln Leu Trp Leu Val Ser Asp Tyr His Glu His Gly Ser Leu  
 275 280 285  
 Phe Asp Tyr Leu Asn Arg Tyr Thr Val Thr Val Glu Gly Met Ile Lys  
 290 295 300  
 Leu Ala Leu Ser Thr Ala Ser Gly Leu Ala His Leu His Met Glu Ile  
 305 310 315 320  
 Val Gly Thr Gln Gly Lys Pro Ala Ile Ala His Arg Asp Leu Lys Ser  
 325 330 335  
 Lys Asn Ile Leu Val Lys Lys Asn Gly Thr Cys Cys Ile Ala Asp Leu  
 340 345 350

Gly Leu Ala Val Arg His Asp Ser Ala Thr Asp Thr Ile Asp Ile Ala  
 355 360 365  
 Pro Asn His Arg Val Gly Thr Lys Arg Tyr Met Ala Pro Glu Val Leu  
 370 375 380  
 Asp Asp Ser Ile Asn Met Lys His Phe Glu Ser Phe Lys Arg Ala Asp  
 385 390 395 400  
 Ile Tyr Ala Met Gly Leu Val Phe Trp Glu Ile Ala Arg Arg Cys Ser  
 405 410 415  
 Ile Gly Gly Ile His Glu Asp Tyr Gln Leu Pro Tyr Tyr Asp Leu Val  
 420 425 430  
 Pro Ser Asp Pro Ser Val Glu Glu Met Arg Lys Val Val Cys Glu Gln  
 435 440 445  
 Lys Leu Arg Pro Asn Ile Pro Asn Arg Trp Gln Ser Cys Glu Ala Leu  
 450 455 460  
 Arg Val Met Ala Lys Ile Met Arg Glu Cys Trp Tyr Ala Asn Gly Ala  
 465 470 475 480  
 Ala Arg Leu Thr Ala Leu Arg Ile Lys Lys Thr Leu Ser Gln Leu Ser  
 485 490 495  
 Gln Gln Glu Gly Ile Lys Met  
 500

## (2) INFORMATION FOR SEQ ID NO:11:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 341 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (vii) IMMEDIATE SOURCE:

- (B) CLONE: KDA-B5

## (ix) FEATURE:

- (A) NAME/KEY: CDS  
 (B) LOCATION: 25..318

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

GGATCCGAAT ACGTGGCGGT TAAA ATA TTC TCC TCC AGG GAT GAG AGA TCT	51
Ile Phe Ser Ser Arg Asp Glu Arg Ser	
1 5	
TGG TTC CGT GAG GCG GAA ATT TAT CAG ACG GTG ATG CTG AGA CAC GAG	99
Trp Phe Arg Glu Ala Glu Ile Tyr Gln Thr Val Met Leu Arg His Glu	
10 15 20 25	
AAC ATC CTC GGT TTC ATC GCA GCT GAC AAC AAA GAT AAT GGA ACT TGG	147
Asn Ile Leu Gly Phe Ile Ala Ala Asp Asn Lys Asp Asn Gly Thr Trp	
30 35 40	
ACA CAA CTC TGG CTT GTG TCA GAG TAT CAC GAG CAG GGC TCC TTG TAT	195
Thr Gln Leu Trp Leu Val Ser Glu Tyr His Glu Gln Gly Ser Leu Tyr	
45 50 55	

GAC TAT TTG AAC AGA AAC ATA GTG ACT GTG GCT GGA ATG GTC AAG CTG 243  
 Asp Tyr Leu Asn Arg Asn Ile Val Thr Val Ala Gly Met Val Lys Leu  
           60                                65                                70

GCG CTT TCC ATA GCG AGT GGT CTG GCT CAC CTG CAC ATG GAG ATC GTG 291  
 Ala Leu Ser Ile Ala Ser Gly Leu Ala His Leu His Met Glu Ile Val  
           75                                80                                85

GGT ACT CAA GGT AAG CTT GCT ATT GCT CACGGTGATA TCAAAAGTCT 338  
 Gly Thr Gln Gly Lys Leu Ala Ile Ala  
           90                                95

AGA --- 341

## (2) INFORMATION FOR SEQ ID NO:12:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 98 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Ile Phe Ser Ser Arg Asp Glu Arg Ser Trp Phe Arg Glu Ala Glu Ile  
   1                                5                                10                                15

Tyr Gln Thr Val Met Leu Arg His Glu Asn Ile Leu Gly Phe Ile Ala  
           20                                25                                30

Ala Asp Asn Lys Asp Asn Gly Thr Trp Thr Gln Leu Trp Leu Val Ser  
           35                                40                                45

Glu Tyr His Glu Gln Gly Ser Leu Tyr Asp Tyr Leu Asn Arg Asn Ile  
           50                                55                                60

Val Thr Val Ala Gly Met Val Lys Leu Ala Leu Ser Ile Ala Ser Gly  
           65                                70                                75                                80

Leu Ala His Leu His Met Glu Ile Val Gly Thr Gln Gly Lys Leu Ala  
           85                                90                                95

Ile Ala

## (2) INFORMATION FOR SEQ ID NO:13:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (vii) IMMEDIATE SOURCE:

- (B) CLONE: PRIMER A

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GCGGATCCGA RTAYGTNGCN GTNAAR

26

## (2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 27 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- (vii) IMMEDIATE SOURCE:
  - (A) LIBRARY: PRIMER B

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

GACTGTAGAR CTYTTDATRT CYCTRTG

27

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 27 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: PRIMER C

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

GACTCTAGAR CTYTTDATRT CNCGR TG

27

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 26 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: PRIMER D

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

GACTCTAGNG AYT T DATRTC YC TRTG

26

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 27 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- (vii) IMMEDIATE SOURCE:

(B) CLONE: PRIMER E

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GACTCTAGAN GAYTTDATRT CNCGRTG

27

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vii) IMMEDIATE SOURCE:

(B) CLONE: PEPTIDE SEQUENCE OF KDA-B5 USED TO DESIGN  
PRIMER A

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Asn Glu Tyr Val Ala Val Lys  
1 5

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 6 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vii) IMMEDIATE SOURCE:

(B) CLONE: PEPTIDE SEQUENCE OF KDA-B5 USED TO DESIGN  
PRIMERS B THRU E

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

His Arg Asp Ile Lys Ser  
1 5



What is claimed is:

1. An isolated DNA molecule comprising a DNA sequence encoding a BMP receptor protein.

5 2. The DNA molecule of claim 1, wherein said DNA sequence is selected from the group consisting of:

(a) nucleotides 61 to 1656 of SEQ ID NO: 1;

(b) nucleotides 247 to 1752 of SEQ ID NO: 3; and

10 (c) sequences which hybridize to (a) or (b) under stringent hybridization conditions and encode a BMP receptor protein with the ability to bind to BMP in a binding assay.

3. An isolated DNA molecule comprising a DNA sequence encoding a serine/threonine kinase receptor protein, wherein the DNA sequence is selected from the group consisting of:

(a) nucleotides 474 to 2000 of SEQ ID NO: 5;

15 (b) nucleotides 80 to 1594 of SEQ ID NO: 7;

(c) nucleotides 83 to 1591 of SEQ ID NO: 9;

(d) sequences which hybridize to any of (c) to (e) under stringent hybridization conditions and encode a serine/threonine receptor protein.

20 4. The DNA molecule of claim 1, wherein said DNA sequence is selected from the group consisting of:

(a) nucleotides encoding for amino acids 24 to 532 of SEQ ID NO: 2; and

(b) nucleotides encoding for amino acids 8 to 502 of SEQ ID NO: 4; and

25 (c) sequences which hybridize to any of (a) or (b) under stringent hybridization conditions and encode a BMP receptor protein with the ability to bind to BMP in a binding assay.

5. An isolated DNA molecule comprising a DNA sequence encoding a serine/kinase receptor protein, wherein the DNA sequence is selected from the group consisting of:

(a) nucleotides encoding for amino acids 18 to 509 of SEQ ID NO: 6; and

30 (b) nucleotides encoding for amino acids 24 to 505 of SEQ ID NO: 8;

and

(c) nucleotides encoding for amino acids 30 to 503 of SEQ ID NO: 10; and

(d) sequences which hybridize to any of (a) to (c) under stringent hybridization conditions and encode a serine/threonine kinase receptor protein.

6. A host cell transformed with the DNA molecule of claim 1.

7. A host cell transformed with the DNA molecule of claim 2.

5 8. A host cell transformed with the DNA molecule of claim 4.

9. A host cell transformed with the DNA molecule of claim 5.

10. An isolated DNA molecule having a sequence encoding a truncated BMP receptor protein which is characterized by the ability to bind to BMP in a binding assay, said DNA molecule comprising a DNA sequence selected from the group consisting of:

(a) nucleotide 61 to 507 of SEQ ID NO:1;

(b) nucleotides encoding amino acids 1 to 149 of SEQ ID NO:2;

(c) nucleotide 247 to 618 of SEQ ID NO:3;

(d) nucleotides encoding amino acids 8 to 124 of SEQ ID NO:4; and

15 (e) naturally occurring allelic sequences and equivalent degenerative codon sequences of (a), (b), (c) or (d).

11. A host cell transformed with the DNA molecule of claim 10.

12. A vector comprising a DNA molecule of claim 10 in operative association with an expression control sequence therefor.

20 13. A method for producing a purified truncated BMP receptor protein, said method comprising the steps of:

(a) culturing in a culture medium a host cell transformed with a DNA sequence according to claim 10, comprising a nucleotide sequence encoding a truncated BMP receptor protein; and

25 (b) recovering and purifying said truncated BMP receptor protein from the culture medium.

14. A purified truncated BMP receptor protein comprising the amino acid sequence from amino acid 24 to amino acid 149 of SEQ ID NO: 2.

30 15. A purified truncated BMP receptor protein comprising the amino acid sequence from amino acid 8 to amino acid 124 of SEQ ID NO: 4.

16. A purified truncated BMP receptor protein produced by the steps of:

(a) culturing in a culture medium a cell transformed with a DNA comprising the nucleotide sequence from nucleotide 61 to 507 of SEQ ID NO: 1; and

5 (b) recovering and purifying from said culture medium a protein comprising the amino acid sequence from amino acid 24 to amino acid 149 of a sequence selected from the group consisting of SEQ ID NO: 2.

17. A purified truncated BMP receptor protein produced by the steps of:

10 (a) culturing in a culture medium a cell transformed with a DNA comprising the nucleotide sequence from nucleotide 247 to 618 of SEQ ID NO: 3; and

(b) recovering and purifying from said culture medium a protein comprising the amino acid sequence from amino acid 8 to amino acid 124 of a sequence selected from the group consisting of SEQ ID NO: 4.

15 18. A method for producing a truncated BMP receptor protein, said method comprising the steps of:

(a) culturing in a culture medium a host cell transformed with a DNA sequence encoding a truncated BMP receptor protein, comprising a truncated ~~nucleotide sequence encoding the ligand binding domain of a BMP receptor~~ protein; and

20 (b) recovering and purifying said BMP receptor protein from the culture medium.

19. A method for producing a truncated BMP receptor protein, said method comprising the steps of:

25 (a) culturing in a culture medium a host cell according to claim 11, comprising a truncated nucleotide sequence encoding the ligand binding domain of a BMP receptor protein; and

(b) recovering and purifying said BMP receptor protein from the culture medium.

30 20. An isolated DNA molecule having a sequence encoding a truncated serine/threonine kinase receptor protein, said DNA molecule comprising a DNA sequence selected from the group consisting of:

(a) nucleotide 474 to 836 of SEQ ID NO:5;

(b) nucleotides encoding amino acids 1 to 121 of SEQ ID NO:6;

(c) nucleotide 80 to 445 of SEQ ID NO:7;

(d) nucleotides encoding amino acids 1 to 122 of SEQ ID NO:8;

(e) nucleotide 83 to 445 of SEQ ID NO:9;

5 (f) nucleotides encoding amino acids 1 to 121 of SEQ ID NO:10; and

(e) naturally occurring allelic sequences and equivalent degenerative codon sequences of (a) through (f).

21. A host cell transformed with the DNA molecule of claim 20.

22. A vector comprising a DNA molecule of claim 20 in operative  
10 association with an expression control sequence therefor.

23. A method for producing a purified truncated serine/threonine kinase receptor protein, said method comprising the steps of:

(a) culturing in a culture medium a host cell transformed with a DNA sequence according to claim 20, comprising a nucleotide sequence encoding a  
15 truncated serine/threonine kinase receptor protein; and

(b) recovering and purifying said truncated serine/threonine kinase receptor protein from the culture medium.

24. A purified truncated serine/threonine kinase receptor protein comprising the amino acid sequence from amino acid 1 to amino acid 121 of  
20 SEQ ID NO: 6.

25. A purified truncated serine/threonine kinase receptor protein comprising the amino acid sequence from amino acid 1 to amino acid 122 of SEQ ID NO: 8.

26. A purified truncated serine/threonine kinase receptor protein comprising the amino acid sequence from amino acid 1 to amino acid 121 of  
25 SEQ ID NO: 10.

27. A method for producing a truncated serine/threonine kinase receptor protein, said method comprising the steps of:

(a) culturing in a culture medium a host cell transformed with a DNA  
30 sequence encoding a truncated serine/threonine kinase receptor protein, comprising a truncated nucleotide sequence encoding the ligand binding domain of a serine/threonine kinase receptor protein; and

(b) recovering and purifying said truncated serine/threonine kinase receptor protein from the culture medium.

28. A method for producing a truncated serine/threonine kinase receptor protein, said method comprising the steps of:

5 (a) culturing in a culture medium a host cell according to claim 21, comprising a truncated nucleotide sequence encoding the ligand binding domain of a serine/threonine kinase receptor protein; and

(b) recovering and purifying said truncated serine/threonine kinase receptor protein from the culture medium.

10 29. A DNA molecule encoding a BMP receptor protein isolated through a method comprising:

(a) preparing a DNA fragment encoding a ligand binding domain of a known BMP receptor protein as a probe;

15 (b) screening a genomic or cDNA library, using the DNA fragment of (a) as a probe;

(c) isolating a DNA molecule which hybridized to the probe identified in step (b); and

~~(d) cloning a DNA molecule encoding a BMP receptor protein from the~~  
DNA molecule of step (c).

20 30. The BMP receptor protein of claim 29, wherein the known BMP receptor protein is CFK1-23a or CFK1-43a.

31. An isolated DNA molecule comprising the DNA sequence of CFK1-23a.

25 32. An isolated DNA molecule comprising the DNA sequence of CFK1-43a.

33. An isolated DNA molecule comprising the DNA sequence of CFK1-10a.

34. An isolated DNA molecule comprising the DNA sequence of W-101.

35. An isolated DNA molecule comprising the DNA sequence of W-120.

30 36. A composition comprising cells transformed with a DNA molecule comprising a DNA sequence encoding one or more BMP receptor proteins.



37. A ligand which binds to a truncated BMP receptor protein, said ligand being selected from the group consisting of an BMP-like monoclonal antibody, a small peptide BMP analogue, and a small organic molecule BMP analogue.

5           38. The ligand of claim 37, comprising a BMP-like monoclonal antibody.

39. The ligand of claim 37, comprising a small peptide BMP analogue.

40. The ligand of claim 37, comprising a small organic molecule BMP analogue.

10           41. A pharmaceutical composition comprising a compound first identified for such use as a ligand for the truncated BMP receptor of claim 20.

42. The composition of claim 41, comprising an BMP-like monoclonal antibody.

43. The composition of claim 41, comprising a small peptide BMP analogue.

15           44. The composition of claim 41, comprising a small organic molecule BMP analogue.

20           45. A ligand for the truncated BMP receptor protein produced by the method of claim 13, said ligand being selected from the group consisting of an BMP-like monoclonal antibody, a small peptide BMP analogue, and a small organic molecule BMP analogue.

46. The ligand of claim 45, comprising an BMP-like monoclonal antibody.

47. The ligand of claim 45, comprising a small peptide BMP analogue.

25           48. The ligand of claim 45, comprising a small organic molecule BMP analogue.

49. A pharmaceutical composition comprising a compound first identified for such use as a ligand for the truncated BMP receptor produced by the method of claim 14.

30           50. The composition of claim 49, comprising an BMP-like monoclonal antibody.

51. The composition of claim 49, comprising a small peptide BMP analogue.

52. The composition of claim 49, comprising a small organic molecule  
BMP analogue.

---

---

---

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C12N15/12 C07K14/71 C07K14/51 C07K16/28 A61K38/18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO,A,94 11502 (LUDWIG INSTITUTE FOR CANCER RESEARCH, GB) 26 May 1994 see the whole document ---	1-36
P,X	WO,A,93 19177 (THE GENERAL HOSPITAL CORP; US) 30 September 1993 see the whole document ---	3,5,9, 20-28
X	DEVELOPMENTAL DYNAMICS, vol.196, no.2, February 1993, US pages 133 - 142 HE, W.H. ET AL.; 'Developmental expression of four novel serine/threonine kinase receptors homologous to the activin/transforming growth factor-beta type II receptor family' see the whole document --- -/--	3,5,9, 20-29

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

- 'A' document defining the general state of the art which is not considered to be of particular relevance
- 'E' earlier document but published on or after the international filing date
- 'L' document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- 'O' document referring to an oral disclosure, use, exhibition or other means
- 'P' document published prior to the international filing date but later than the priority date claimed

- 'T' later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- 'X' document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- 'Y' document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- '&' document member of the same patent family

Date of the actual completion of the international search

20 December 1994

Date of mailing of the international search report

30. 12. 94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

Nauche, S

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,91 18047 (GEENETECH, INC.; US) 28 November 1991 see the whole document ---	37-52
X	WO,A,88 00205 (GENETICS INSTITUTE, INC.; US) 14 January 1988 see the whole document ---	37-52
X	EP,A,0 416 578 (TADEKA CHEMICAL INDUSTRIES, LTD.; JP) 13 March 1991 see the whole document ---	37-52
X	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA., vol.89, December 1992, WASHINGTON US pages 11740 - 11744 CUNNINGHAM, N.S. ET AL.; 'Osteogenin and recombinant bone morphogenetic protein 2B are chemotactic for human monocytes and stimulate transforming growth factor beta1 m RNA expression' see the whole document -----	37-52

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9411502	26-05-94	AU-B- 5432094	08-06-94
WO-A-9319177	30-09-93	AU-B- 3920693	21-10-93
WO-A-9118047	28-11-91	US-A- 5168050	01-12-92
		AT-T- 114163	15-12-94
		EP-A- 0531448	17-03-93
WO-A-8800205	14-01-88	US-A- 4877864	31-10-89
		AU-B- 613314	01-08-91
		AU-A- 7783587	29-01-88
		EP-A- 0313578	03-05-89
		JP-A- 6298800	25-10-94
		JP-T- 2500241	01-02-90
		US-A- 5013649	07-05-91
		US-A- 5166058	24-11-92
		US-A- 5187076	16-02-93
		US-A- 5116738	26-05-92
		US-A- 5366875	22-11-94
		US-A- 5106748	21-04-92
		US-A- 5108922	28-04-92
		US-A- 5141905	25-08-92
EP-A-0416578	13-03-91	JP-A- 4154799	27-05-92



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**